Statistical mechanics model for Clifford random tensor networks and monitored quantum circuits

Yaodong Li¹, Romain Vasseur, Matthew P. A. Fisher, and Andreas W. W. Ludwig¹ Department of Physics, University of California, Santa Barbara, California 93106, USA Department of Physics, University of Massachusetts, Amherst, Massachusetts 01003, USA

(Received 10 October 2023; revised 4 April 2024; accepted 5 April 2024; published 13 May 2024)

We study entanglement transitions in Clifford (stabilizer) random tensor networks (RTNs) and monitored quantum circuits, by introducing an exact mapping onto a (replica) statistical mechanics model. For RTNs and monitored quantum circuits with random Haar unitary gates, entanglement properties can be computed using statistical mechanics models whose fundamental degrees of freedom ("spins") are permutations, because all operators commuting with the action of the unitaries on a tensor product Hilbert space are (linear combinations of) permutations of the tensor factors ("Schur-Weyl duality"). When the unitary gates are restricted to the smaller group of Clifford unitaries, the set of all operators commuting with this action, called the *commutant*, will be larger, and no longer form a group. We use the recent full characterization of this commutant by Gross et al. [Commun. Math. Phys. 385, 1325 (2021)] to construct statistical mechanics models for both Clifford RTNs and monitored quantum circuits, for on-site Hilbert-space dimensions which are powers of a prime number p. The elements of the commutant form the spin degrees of freedom of these statistical mechanics models, and we show that the Boltzmann weights are invariant under a symmetry group involving orthogonal matrices with entries in the finite number field \mathbf{F}_p ("Galois field") with p elements. This implies that the symmetry group and consequently all universal properties of entanglement transitions in Clifford circuits and RTNs will, respectively, in general depend on and only on the prime p. We show that Clifford monitored circuits with on-site Hilbert-space dimension $d = p^M$ are described by percolation in the limits $d \to \infty$ at (a) p = fixed but $M \to \infty$, and at (b) M = 1 but $p \to \infty$. In the limit (a) we calculate the effective central charge, and in the limit (b) we derive the following *universal* minimal cut entanglement entropy $S_A = (\sqrt{3}/\pi) \ln p \ln L_A$ for d = p large at the transition. We verify those predictions numerically, and present extensive numerical results for critical exponents at the transition in monitored Clifford circuits for prime number on-site Hilbert-space dimension d = p for a variety of different values of p, finding that projective and forced measurement schemes yield the same critical exponents and that they approach percolation values at large p. We clearly establish multifractal scaling of the purity, reflected in a continuous spectrum of critical exponents, while the typical exponent is the prefactor of the logarithm in the entanglement entropy. As a technical result, we generalize the notion of the Weingarten function, previously known for averages involving the Haar measure, to averages over the Clifford group.

DOI: 10.1103/PhysRevB.109.174307

I. INTRODUCTION

Entanglement plays a central role in the physics of closed many-body quantum systems, in both equilibrium and nonequilibrium settings [1–5]. With the advent of noisy intermediate-scale quantum (NISQ) devices [6], the study of the dynamics of quantum information in open quantum systems has attracted a lot of attention recently. The interplay of unitary many-body quantum evolution [7–12]—which generates quantum entanglement—and the nonunitary processes generated by noisy couplings to the environments or by measurements—which tend to reveal and destroy quantum information—leads to a broad variety of new dynamical phases of matter and phase transitions.

A particularly interesting example that captures the competition between unitary dynamics and nonunitary processes is the so-called measurement-induced phase transition [13,14]. A simple setup where such a transition occurs is provided by "monitored" quantum quantum circuits made up of random unitary gates, combined with local projective measurements occurring at a fixed rate. As a function of the measurement probability p_0 , a remarkable entanglement transition occurs. At low p_0 , the unitary dynamics can efficiently scramble quantum information into highly nonlocal degrees of freedom that cannot be accessed by the local measurements [15-18]. In that regime, the system reaches a highly entangled steady state where the entanglement entropy of a subsystem scales as its volume (volume law). In contrast, at high measurement probability p_0 , the frequent local measurements are able to effectively collapse the wave-function of the system, reminiscent of a many-body Zeno effect, with the entanglement of subsystems scaling as their boundary (area law). This fundamentally new transition is not observable in the mixed density matrix of the system averaged over measurement outcomes, but is apparent in individual quantum trajectories of the pure state wave function, conditional on measurement outcomes. This transition has been studied extensively in recent years, generalized to various classes of quantum dynamics with different symmetries and dimensionality [13-54], and realized experimentally in a trapped-ion quantum computer [55].

An apparently different, but closely related entanglement transition was discovered a bit earlier in random tensor

see e.g., [64]. Numerical evidence indicates that the measurement induced transition in Clifford qubit circuits is in a

different universality class from the Haar case, but a theoreti-

cal description of the Clifford transition has remained elusive.

Part of the difficulty is that while the structure of the Clifford

gates allows for efficient classical simulations, the very same

structure makes a general theoretical formulation of averages

networks [33,34,56–58]. There the transition can be induced by tuning the bond dimension¹ of a state obtained at the boundary of two-dimensional random tensor networks previously formulated deep in the volume law phase, far from the transition in Ref. [59]. A unifying description of such entanglement transitions was proposed in Refs. [24,25,56] in terms of a replica statistical mechanics model. A replica trick is needed to deal with the intrinsic nonlinearities of the problem-the entanglement transition is not visible in quantities that are linear in the density matrix of the system. The average over either Gaussian random tensors or Haar unitary gates in the replicated tensor network can then be performed exactly and leads to effective degrees of freedom ("spins") that are permutations of the replicas. There is a deep mathematical reason for the emergence of the permutation group here, which is known as the Schur-Weyl duality: all operators commuting with the action of the unitary group on a tensor product Hilbert space are linear combinations of permutations of the tensor factors (replicas). This leads to an exact reformulation of the problem of computing entanglement entropies and other observables in Haar monitored quantum circuits or random tensor networks in terms of (the replica limit of) a two-dimensional classical statistical mechanics model with permutation degrees of freedom, and ferromagnetic interactions. Even if taking the replica limit explicitly is challenging in general, this mapping ends up explaining most (if not all) of the phenomenology of entanglement transitions, which in that language become simple symmetry-breaking (ordering) transitions. In particular, it naturally explains the emergence of conformal invariance (with a dynamical exponent z = 1) at criticality [19,21,25,56]. The entanglement entropy maps onto the free energy cost of "twisting" the boundary conditions in the entanglement interval, forcing the insertion of a domain wall at the boundary of the entanglement interval. This free-energy cost scales with the size of the interval in the ferromagnetic (symmetry-broken) phase of the statistical mechanics model, corresponding to the volume-law entangled phase, while it remains order one in the paramagnetic phase, corresponding to the area-law phase. Additional simplifications occur in the limit of large on-site Hilbert-space dimensions, where the replica limit can be taken analytically, predicting an entanglement transition that falls into the classical percolation universality class [24,25].

While these analytical results were derived for Haar random unitary gates or tensors, large-scale numerical results are conveniently obtained using random Clifford unitary gates, and single-qubit measurements restricted to the Pauli group. Such Clifford circuits (and the related stabilizer tensor networks [58,60]) can be efficiently simulated on a classical computer in times polynomial in the system size thanks to the Gottesman–Knill theorem [61–63], whereas the simulation times of Haar circuits usually scale exponentially with the system size. Clifford circuits have proven useful in the study of entanglement and operator dynamics in various contexts, over Clifford gates more challenging than over featureless Haar gates. In this paper, we leverage recent results on the "commutant" of the Clifford group [65], which generalize the notion of Schur-Weyl duality to the Clifford case, to derive a replica statistical mechanics description of Clifford monitored quantum circuits and stabilizer random tensor networks for qudits with dimension $d = p^M$, where p is a prime number [66]. The elements of the commutant replace permutations to form the spin degrees of freedom of the resulting statistical mechanics models. We show that the Boltzmann weights are invariant under a symmetry group involving orthogonal matrices with entries in the finite number field \mathbf{F}_p ("Galois field") with p elements. This implies that the symmetry group and all universal properties of entanglement transitions in Clifford circuits and random tensor networks will in general depend on the on-site Hilbert dimension $d = p^M$ of the circuit, but only via the prime p, i.e., they are independent of the power M. This also explains that Clifford circuits and random stabilizer tensor networks (i.e., Clifford random tensor networks) are in different universality classes than their Haar counterparts. In particular, in the latter cases the on-site Hilbert-space dimension is a short-distance feature that does not affect their symmetry [25] and thus not the universality class (as long as d is finite), in contrast with the Clifford cases. Our approach also allows us to derive mappings onto classical percolation in the limit of large on-site Hilbert-space dimension d. Our predictions are supported by exact numerical simulations of large monitored Clifford circuits for d = p up to p = 997, and for $d = p^M$ at a few small values of M, which also show that projective and forced measurement schemes yield the same critical exponents for Clifford circuits. Our approach also explains some recent numerical results on Clifford random tensor networks [58]. Finally, we provide clear evidence that the purity of the reduced density matrix of a finite interval exhibits multifractal scaling at the transition, by numerically investigating the statistical fluctuations of the entanglement entropy about its mean. This implies different scaling exponents for the averaged and the typical purity (the latter being equal to the prefactor of the logarithm of system size in the entanglement entropy) and is reflected in a continuous spectrum of critical exponents associated with purity scaling.

This paper is organized as follows: In Sec. II, we focus on random tensor networks (RTNs) and extend the notion of Schur-Weyl duality to the Clifford case, relying heavily on Ref. [65]. We describe the structure of the commutant and explain how the statistical mechanics model describing Haar RTNs can be extended to Clifford RTNs. In Sec. III, we generalize the notion of Weingarten functions to the Clifford case and apply those results to monitored Clifford circuits. We also make concrete predictions in the limit of large on-site Hilbert space, carefully distinguishing the cases of p or M fixed when $d = p^M \to \infty$. In Sec. IV, we provide numerical evidence that the universality class of the entanglement transition at

¹Or, at a fixed sufficiently large bond dimension, by randomly diluting (=eliminating) bulk bonds with probability p_0 , the latter serving as the parameter tuning through the transition [56].



FIG. 1. (a) A tensor T_v at vertex v with coordination number z = 4. (b) Arranging the tensors in a network (taken to be square lattice here), we obtain a "tensor network state" $\rho[T]$ [see Eq. (4)] by contracting indices on all bulk edges, as implemented by projecting onto the maximally entangled state $|I_e\rangle$. The state $\rho[T]$ is supported on the boundary qudits (highlighted by purple squares) and is in general unnormalized.

qudit dimension $d = p^M$ indeed depends on p, but not on M, report on the equality of exponents for projective and forced measurement schemes, and establish multifractal scaling of the purity. Technical details and additional mathematical results are gathered in the Appendixes.

II. RANDOM TENSOR NETWORKS AND SCHUR-WEYL DUALITY

A. Review of Haar case

Following Ref. [56] we consider a Tensor Network with a set of vertices v, from each of which emerge z edges ve_1, ve_2, \ldots, ve_z , where z is the coordination of the vertex (assumed to be the same for all vertices), see Fig. 1(a). We denote the bond dimension at each edge by d. At each vertex v there is a quantum state described by a tensor

$$|T_{v}\rangle = \sum_{\mu_{ve_{1}},\mu_{ve_{2}},\dots,\mu_{ve_{z}}=1}^{a} (T)_{\mu_{ve_{1}},\mu_{ve_{2}},\dots,\mu_{ve_{z}}}$$
$$\times |\mu_{ve_{1}}\rangle \otimes |\mu_{ve_{2}}\rangle \otimes \dots \otimes |\mu_{ve_{z}}\rangle = \sum_{\mu_{v}=1}^{D} T_{\mu_{v}}|\mu_{v}\rangle,$$
(1)

where we introduced the shorthand $\mu_v = (\mu_{ve_1}, \mu_{ve_2}, \dots, \mu_{ve_z})$ for the collection of labels of the $D := (d)^z$ basis elements at a vertex, $\mu_v = 1, \dots D$. Defining a maximally entangled state on the edge $e := \langle v, v' \rangle$ connecting two adjacent vertices v and v',

$$|I_e\rangle := \frac{1}{\sqrt{d}} \sum_{\mu_{ve}, \mu_{v'e}=1}^d \delta_{\mu_{ve}, \mu_{v'e}} |\mu_{ve}\rangle \otimes |\mu_{v'e}\rangle, \qquad (2)$$

the (unnormalized) wave function of the tensor network is defined by

$$|\psi[T]\rangle := (\otimes_e \langle I_e|)(\otimes_v | T_v \rangle), \tag{3}$$

where the tensor product is over all (bulk) edges and all vertices of the network, respectively. The object of interest is

the (so far still unnormalized) pure state density matrix of the network

$$\rho[T] := |\psi[T]\rangle \langle \psi[T]|, \tag{4}$$

for the remaining (uncontracted) degrees of freedom ("dangling legs" of the tensor network), which we choose to be at the boundary of the network, see Fig. 1(b). Owing to the need to normalize the states of the wave function at the boundary of the network (and to describe the *n*th Rényi entropies for any integer *n*), it is necessary to work with the tensor product of Q = nm copies of this (unnormalized) density matrix:

$$\otimes^{\mathcal{Q}} \rho[T] = \otimes^{\mathcal{Q}}(|\psi[T]\rangle \langle \psi[T]|). \tag{5}$$

More precisely, our goal is to compute the *n*th entanglement Rényi entropy of a subregion *A* of the dangling ("boundary") legs of the tensor networks:

$$S_{A}^{(n)} = \frac{1}{1-n} \ln \frac{\text{tr}\rho_{A}^{n}}{(\text{tr}\rho)^{n}},$$
(6)

with $\rho_A = \text{tr}_{\bar{A}}\rho$ being the reduced density matrix in the region *A*, and \bar{A} the complement of *A*. The average of this quantity over the random tensors, to be defined more precisely below, can be computed from the replicated density matrix using

$$S_{A}^{(n)} = \frac{1}{1 - n} \lim_{m \to 0} \frac{1}{m} \Big[\Big(\mathrm{tr} \rho_{A}^{n} \Big)^{m} - (\mathrm{tr} \rho^{\otimes Q}) \Big], \tag{7}$$

where $(\operatorname{tr} \rho_A^n)^m$ and $\operatorname{tr} (\rho^{\otimes Q})$ differ only in the way the boundary legs are contracted.

The Haar random tensor network is defined by letting the state $|\mu_v^{(i)}\rangle$ at each vertex v in each (replica and Rényi) copy i = 1, ..., Q arise from a fixed basis state in the *D*dimensional Hilbert space associated with the vertex by the action of the same unitary $D \times D$ matrix² U in each (replica and Rényi) copy, chosen randomly and independently at each vertex v from the Haar ensemble. Explicitly, the tensor in (1) will then be of the form

$$T_{\mu_v} = \langle \mu_v | \hat{U} | 1 \rangle \coloneqq (U)_{\mu_v, 1}, \quad \mu_v = 1, \dots, D.$$
(8)

The tensor factor contribution to (5) arising from the quantum states (1) at vertex v will then be, using (3),

$$\begin{split} \otimes^{Q}(|T_{v}\rangle \langle T_{v}|) &= \otimes^{Q}(\hat{U}|1\rangle \langle 1|\hat{U}^{\dagger}) \\ &= \sum_{\mu_{v}^{(i)}, v_{v}^{(j)}} U_{\mu_{v}^{(1)}, 1} \cdots U_{\mu_{v}^{(Q)}, 1} U^{\dagger}_{1, v_{v}^{(1)}} \cdots U^{\dagger}_{1, v_{v}^{(Q)}} \\ &\times (|v_{v}^{(1)}\rangle \otimes \cdots \otimes |v_{v}^{(Q)}\rangle) \otimes (\langle \mu_{v}^{(1)}| \otimes \cdots \otimes \langle \mu_{v}^{(Q)}|). \end{split}$$

$$(9)$$

Specializing the familiar result³ for the average $\mathbb{E}_{U \in U(D)}[\ldots]$ over the Haar measure $d\mu(U)$ [67]

$$\int_{\text{Haar}} d\mu(U) U_{i_1, j_1} \cdots U_{i_Q, j_Q} U_{j'_1, i'_1}^{\dagger} \cdots U_{j'_Q, i'_Q}^{\dagger}$$
$$= \sum_{\sigma, \tau \in S_Q} Wg(\tau \sigma^{-1}) \delta_{i_1, i'_{\sigma(1)}} \cdots \delta_{i_Q, i'_{\sigma(Q)}} \delta_{j_1, j'_{\tau(1)}} \cdots \delta_{j_Q, j'_{\tau(Q)}}$$
(10)

³A derivation is also provided in (C5) of Appendix C.

²Recall that *d* is the bond dimension of the Hilbert space on each edge, and $D = d^z$ (z = coordination number) is the dimension of the total Hilbert space associated with each vertex *v*.

to the case where $j_1 = \cdots = j_Q = 1$ and $j'_1 = \cdots = j'_Q = 1$ yields

$$\mathbb{E}_{U \in U(D)} \Big[U_{i_1,1} \cdots U_{i_Q,1} U_{1,i'_1}^{\dagger} \cdots U_{1,i'_Q}^{\dagger} \Big]$$
$$= \frac{1}{\mathcal{N}_H} \sum_{\tau \in S_Q} \delta_{i_1,i'_{\tau(1)}} \cdots \delta_{i_Q,i'_{\tau(Q)}},$$

where $1/\mathcal{N}_H = \sum_{\sigma} Wg_D(\sigma) = \prod_{k=1}^{Q} (D+k-1)^{-1}$. This implies that the Haar average of (9) gives⁴ a contribution coming from the vertex v of the form

$$\mathbb{E}_{U \in U(D)}[\otimes^{Q}(|T_{v}\rangle\langle T_{v}|)] = \frac{1}{\mathcal{N}_{H}} \sum_{\tau \in S_{Q}} \hat{R}_{v}(\tau), \qquad (11)$$

where

$$\hat{R}_{v}(\tau) := \sum_{\vec{\mu}_{v}} |\hat{\tau}(\vec{\mu}_{v})\rangle \langle \vec{\mu}_{v}|, \quad \tau \in S_{Q},$$
(12)

and where $\vec{\mu}_v = (\mu_v^{(1)}, \mu_v^{(2)}, \dots, \mu_v^{(Q)})$ labels an orthonormal basis of the *Q* (replica and Rényi) copies of the *D*-dimensional Hilbert space, $(\mathbf{C}^D)^Q$, at vertex *v*. Here S_Q denotes the permutation group of *Q* elements, and $\hat{\tau}$ is the operator that permutes the *Q* factors of the tensor product,

$$|\vec{\mu}_{v}\rangle := |\mu_{v}^{(1)}, \mu_{v}^{(2)}, \dots, \mu_{v}^{(Q)}\rangle, |\hat{\tau}(\vec{\mu}_{v})\rangle := |\mu_{v}^{(\tau(1))}, \mu_{v}^{(\tau(2))}, \dots, \mu_{v}^{(\tau(Q))}\rangle.$$
(13)

The operator $\hat{R}_v(\tau)$ in (12) is simply the permutation operator permuting the Q (replica and Rényi) copies of the vector space \mathbf{C}^D at vertex v. Equation (11) is the key result for the Haar case. The operator $\hat{R}_v(\tau)$ in (12) commutes with the action of all unitaries $U \in U(D)$ on the tensor product $(\mathbf{C}^D)^Q$ associated with a vertex. Viewing the set of operators $\hat{R}(\tau)$ where τ runs over all permutations in the group S_Q as a basis of a (complex) linear vector space,⁵ it can be shown that this vector space of operators forms the set of all operators commuting with the action of the unitaries on the tensor product $(\mathbf{C}^D)^Q$. This set of operators is called the *commutant* of action of the unitary group on the tensor product, and the stated result is known as the famous statement of *Schur-Weyl duality*.

Consider now a single edge *e* emanating from a vertex *v*. We label an orthonormal basis of the *Q* (replica and Rényi) copies of the *d*-dimensional Hilbert space of edge *e* at vertex *v* by $\vec{\mu}_{ve} = (\mu_{ve}^{(1)}, \mu_{ve}^{(2)}, \dots, \mu_{ve}^{(Q)})$, corresponding to the kets

$$|\vec{\mu}_{ve}\rangle := \left|\mu_{ve}^{(1)}, \mu_{ve}^{(2)}, \dots, \mu_{ve}^{(Q)}\right\rangle.$$
 (14)

Recalling the decomposition from (1),

$$|\mu_{v}\rangle = |\mu_{ve_{1}}\rangle \otimes |\mu_{ve_{2}}\rangle \otimes \cdots \otimes |\mu_{ve_{z}}\rangle, \tag{15}$$

valid in each (replica and Rényi) copy, as well as (13), we see that the operator in (12) is the tensor product of operators

$$\hat{R}_e(\tau),\tag{16}$$

⁵Which is in fact an algebra because we can multiply the operators.

defined on the edges e of the vertex v, where

$$\hat{R}_{e}(\tau) := \sum_{\vec{\mu}_{ve}} |\hat{\tau}(\vec{\mu}_{ve})\rangle \langle \vec{\mu}_{ve}|, \quad \tau \in S_{Q},$$
(17)

and $\hat{\tau}$ labels⁶ the operator that permutes the Q (replica and Rényi) copies of the *d*-dimensional Hilbert space, $(\mathbf{C}^d)^Q$, at an edge *e* at vertex *v*:

$$|\vec{\mu}_{ve}\rangle := |\mu_{ve}^{(1)}, \mu_{ve}^{(2)}, \dots, \mu_{ve}^{(Q)}\rangle, |\hat{\tau}(\vec{\mu}_{ve})\rangle := |\mu_{ve}^{(\tau(1))}, \mu_{ve}^{(\tau(2))}, \dots, \mu_{ve}^{(\tau(Q))}\rangle.$$
(18)

In other words, we have

$$\hat{R}_{v}(\tau) = \bigotimes_{e=1}^{z} \hat{R}_{e}(\tau) = [\hat{R}_{e}(\tau)]^{\otimes z}.$$
(19)

Consider now two adjacent vertices v and v' connected by an edge e. The contribution to (5) arising from the edge joining adjacent vertices v (permutation τ) and v' (permutation σ) is then from (3) and (5)

$$(\otimes^{\mathcal{Q}}\langle I_e|)R_e(\tau)R_e(\sigma)(\otimes^{\mathcal{Q}}|I_e\rangle)$$

$$= \frac{1}{d^{\mathcal{Q}}}\sum_{\vec{\mu}_{ve},\vec{\mu}_{v'e}} [\hat{R}_e(\tau)]_{\vec{\mu}_{ve},\vec{\mu}_{v'e}} [\hat{R}_e(\sigma)]_{\vec{\mu}_{v'e},\vec{\mu}_{ve}}$$

$$= \frac{1}{d^{\mathcal{Q}}} \operatorname{tr}[\hat{R}_e(\tau)\hat{R}_e^{\dagger}(\sigma)] = \frac{1}{d^{\mathcal{Q}}} \operatorname{tr}[\hat{R}_e(\tau\sigma^{-1})]. \quad (20)$$

It is seen by inspection [56] [using definition (17)] that

$$\operatorname{tr}[\hat{R}_{e}(\tau\sigma^{-1})] = e^{\ln(d)C(\tau\sigma^{-1})}, \qquad (21)$$

where $C(\tau)$ denotes the number of cycles in the permutation $\tau \in S_Q$. Once the product over the analogous contributions from all edges and vertices is taken, a sum over all independent permutations σ_v in the group S_Q at all vertices v has to be performed. This yields the (bulk) partition function of the Haar stat mech model [56], whose spins are the elements of the permutation group located at the vertices,

$$Z_{\emptyset} = \mathbb{E}_{U \in U(D)} \operatorname{tr} \rho^{\otimes Q} = \prod_{\{\sigma_v\}} \exp\left\{-\sum_{\langle v, v' \rangle} \mathcal{H}(\sigma_v, \sigma_{v'})\right\},$$
$$\mathcal{H}(\sigma_v, \sigma_{v'}) = \ln(d) |\sigma_v, \sigma_{v'}| := \ln(d) [Q - C(\sigma_v \sigma_{v'}^{-1})],$$
(22)

where we can think of $|\sigma_v, \sigma_v|$, defined above, as a "metric" on permutations.—We now see that the fact that the degrees of freedom (the spins) of the RTN model of random Haar tensors are the elements of the permutation group arises from the fact that the permutations form a basis of the commutant of the action of the unitaries on the tensor product Hilbert space. The permutation degrees of freedom at the end of the dangling boundary legs are fixed and are dictated by the trace under consideration. To compute $Z_{\emptyset} = \mathbb{E}_{U \in U(D)} \operatorname{tr} \rho^{\otimes Q}$, the boundary permutations are fixed to the identity, corresponding to contracting each replica with itself. On the other hand, one can compute $Z_A = \mathbb{E}_{U \in U(D)}(\operatorname{tr} \rho_A^n)^m$ by fixing the same

⁴The sum over permutations τ was replaced by the sum over τ^{-1} , which makes no difference.

⁶Upon some slight abuse of notation.

identity permutation in the region \overline{A} , while fixing $g_{SWAP} = (1, 2, ..., n)^{\otimes m}$ in the region A in order to compute the partial trace.

In the sequel, we identify, using the same logic, the degrees of freedom (i.e., the spins) of the RTN model for random Clifford tensors [58,60]. Analogously, they will form a basis of the commutant of the action of the Clifford unitaries on the Q-fold tensor product. Since Clifford unitary tensors form a subgroup of all (Haar) unitary tensors, the set of operators that will commute with their action on the tensor Hilbert space will be larger. In other words, the commutant will be larger as compared with the Haar case. We now proceed to explain the nature of this larger commutant, and thus of the spins of the stat mech model for the RTN model of random Clifford tensors.

B. Schur-Weyl duality for the Clifford random tensor network model

Throughout this section (and any discussion of the RTN or monitored quantum circuits with random Clifford unitaries in this paper) we take the on-site Hilbert-space dimension d at each edge to be a power of a prime number p, i.e., $d = p^M$ with some integer M.

The difference with the case of Haar unitary tensors appears in (8): In the Clifford case the matrix U is replaced by a unitary matrix, which we denote by V, in the Clifford subgroup Cliff(N, p) of all unitary $D \times D$ matrices U(D), where $D = d^z = p^{zM} = p^N$, and N = zM. First, we use this setup to describe the commutant of this action on the Q-fold tensor product space. Second, after the description of the commutant is completed, we consider the generalization of the average (11) to an average over all the elements of this Clifford subgroup Cliff(N, p) acting on a fixed tensor $|T_v\rangle$ which is now chosen to be a stabilizer state.

For the first step, the key statement of Schur-Weyl duality for the Clifford group is that there is a specific generalization of the operators (12) spanning the commutant, which are no longer characterized by permutations.

1. Description of properties of the commutant

Specifically, we are interested in the basis $|\vec{\mu}_v\rangle$ of the *Q*-fold tensor product space $(\mathbf{C}^D)^Q$ of all *z* edges⁷ emerging from a given vertex *v* so that we can write $(\mathbf{C}^D)^Q = [(\mathbf{C}^p)^{\otimes N}]^{\otimes Q}$, where N = Mz. The *p* basis elements of the vector space \mathbf{C}^p can be labeled by the *p* elements of $\mathbf{Z}_p := \mathbf{Z}/(p \cdot \mathbf{Z}) = 0, 1, 2, ..., (p-1)$, i.e., by the set of integers modulo *p* (the "computational basis"). It will soon become important that, when *p* is a prime, the finite set \mathbf{Z}_p forms a number field.⁸ which is often also denoted by $\mathbf{F}_p = \mathbf{Z}_p$.

Let us first consider the simplest case where N = 1, i.e., of the *Q*-fold tensor product $(\mathbb{C}^p)^{\otimes Q}$, which turns out to be the most important case, underlying all others. The same unitary $p \times p$ matrix *V* in the Clifford subgroup Cliff(1, p) of the unitary group U(p) acts simultaneously on each of the *Q* factors of the tensor product, generalizing (8) with *D* replaced by *p*. Since the basis elements of \mathbf{C}^p are labeled (as above) by elements $\mu \in \mathbf{Z}_p = \mathbf{F}_p$, the elements of a basis of $(\mathbf{C}^p)^{\otimes Q}$ will be labeled by *Q*-dimensional (say, column-) vectors⁹ with entries in $\mathbf{Z}_p = \mathbf{F}_p$, written as usual as elements of $(\mathbf{Z}_p)^Q = (\mathbf{F}_p)^Q$, i.e., by vectors $\vec{\mu} = (\mu^{(1)}, \mu^{(2)}, \dots, \mu^{(Q)})^t$, where $\mu^{(1)}, \dots, \mu^{(Q)} \in \{0, 1, \dots, (p-1)\} = \mathbf{F}_p$. In other words, every *Q*-dimensional (column) vector with entries in $\mathbf{Z}_p = \mathbf{F}_p$, i.e., every (column) vector in $(\mathbf{Z}_p)^Q = (\mathbf{F}_p)^Q$, denotes a basis element of the vector space $(\mathbf{C}^p)^Q$.

The main result of Ref. [65] is that, in this case, the generalization of (12) and (17), i.e., of the operators which in the Haar case formed a basis of the commutant, i.e., of the set (actually the vector space) of all operators commuting with the action of the unitary group on the tensor product, in the Clifford case is based on the operators

$$\hat{r}(T) = \sum_{(\vec{v},\vec{\mu})\in T} |\vec{v}\rangle\langle\vec{\mu}|, \qquad (23)$$

where *T* runs over a certain set (to be specified below) of subspaces of $(\mathbf{Z}_p)^Q \oplus (\mathbf{Z}_p)^Q$. In other words, the permutation in the Haar case is now replaced by the (more general) object *T*. The so-defined $\hat{r}(T)$ is clearly an operator acting on $(\mathbf{C}^p)^Q$. The set of allowed subspaces *T*, denoted by $\Sigma_Q(p)$ and to be specified below, forms the commutant in the Clifford case.

To appreciate the meaning of (23), let us first look at a special case: Since the set of all Clifford unitary matrices $V \in \text{Cliff}(1, p)$ forms a subgroup of the set of all unitary matrices $U \in U(p)$, the set of all operators commuting with the action of all *Clifford* unitaries on the *Q*-fold tensor product of the Hilbert space \mathbb{C}^p must be a larger set than the set of all those commuting with the action of *all* unitaries *U*. Since we know from the previous section on the Haar case that the commutant in the latter case is spanned by all permutations $\tau \in S_Q$, the operators that appeared in (12) and (17) must form a subset of the set of operators described in (23). Indeed, choosing the subspace of $(\mathbb{Z}_p)^Q \oplus (\mathbb{Z}_p)^Q$ to be of the *special* form

$$T = \{ (\hat{\tau}(\vec{\mu}), \vec{\mu}) : \vec{\mu} \in (\mathbf{Z}_p)^Q \}$$
(24)

reproduces precisely the previous expression from (12) and (17), where in the above equation $\hat{\tau}$ acts by permuting the Q rows of the column vector $\vec{\mu}$.

Before we move on to the description of the commutant and its properties for the Clifford case, we state an important *stabilization property* [65] that says that, once the prime number *p* is fixed, this commutant does not depend on the dimension of the (on-site) Hilbert space on which it acts, as long as that dimension is large enough. This is a very important statement because it says that, for a fixed prime *p*, the specific form of the commutant that we describe below is universally valid, independent of the (sufficiently large) prime-power dimension of Hilbert space it acts on. For this purpose, we consider a general on-site Hilbert space (\mathbb{C}^p)^N of dimension $D = p^N$ (as above), i.e., simply the *N*-fold tensor product of the Hilbert space \mathbb{C}^p discussed above. We now consider as

 $^{^{7}}$ As in (13).

⁸Meaning that addition, multiplication, and division are defined in the usual way.

⁹The superscript ^{*t*} denotes the transpose.

before the tensor product of Q (replica, and Rényi) copies of this *D*-dimensional Hilbert space, i.e., $[(\mathbf{C}^p)^N]^{\otimes Q} = (\mathbf{C}^p)^{N \cdot Q}$, and on each of the Q copies of the $D = p^N$ -dimensional vector space $(\mathbf{C}^p)^N$ acts the same unitary $D \times D$ matrix V in the Clifford subgroup Cliff(N, p) of the unitary group U(D). Now the following statement holds:

Stabilization property [65]. The number of linearly independent operators (which will span the commutant) that commute with the action of the same Clifford unitary matrices $V \in \text{Cliff}(N, p)$ acting simultaneously on each of the Qfactors of the Q-fold tensor product, $[(\mathbf{C})^D]^{\otimes Q}$, where $D = p^N$ (p = prime), is *independent of* N as long as $N \ge (Q - 1)$.

Moreover, these linearly independent operators are simply

$$\hat{R}(T) = [\hat{r}(T)]^{\otimes N}$$
, where $T \in \Sigma_Q(p)$, (25)

i.e., they are nothing but the *N*-fold tensor products of the operators introduced in (23) above. Note that these operators are thus uniquely determined by the set $\Sigma_Q(p)$, independent of the power *N* determining the Hilbert-space dimension $D = p^N$, as long as *N* is large enough¹⁰ (as specified above).

The key result of Ref. [65] is a complete characterization of the set $\Sigma_Q(p)$ of subspaces *T* appearing in (25), which describes the commutant of the Clifford action on the *Q*-fold tensor product. A more detailed summary of this is given in Appendix A. Here we just list those properties of the commutant $\Sigma_Q(p)$ that appear directly in the formulation of the Clifford RTN model. These are a metric on the commutant, and the invariance of this metric under a certain symmetry group. We now proceed to discuss these two properties.

An important property of the commutant $\Sigma_Q(p)$ is that it contains a group as a subset [though $\Sigma_O(p)$ is in general larger than this group and does in general not form a group]. This group is the group of all orthogonal $Q \times Q$ matrices *O* with elements in the number field $\mathbf{Z}_p = \mathbf{F}_p$ which satisfy besides the "orthogonality condition" $O^t O = OO^t = I_Q$ (the $Q \times Q$ identity matrix) also the additional condition of "being stochastic," meaning that the column vector 1_0 whose entries are all the number $1 \in \mathbf{F}_p$, is invariant under the action of the matrix O, as well as of its transpose, i.e., $O 1_Q = O^t 1_Q = 1_Q$. The set of such matrices forms a group denoted by $\mathcal{O}_Q(p)$, the so-called stochastic orthogonal group. Note that [in connection with the discussion around (24)] the permutation group S_Q is a subgroup of $\mathcal{O}_Q(p)$, namely, it corresponds to the set of those $Q \times Q$ matrices which have only one $1 \in \mathbf{F}_p$ in each row and each column; those matrices clearly implement permutations.

For the formulation of the Clifford stat mech model we use the fact that there exists a *metric* on the commutant: Namely, that for any two elements T_a and T_b of the commutant $\Sigma_Q(p)$ there exists a *metric* which arises from the trace

$$W(T_a, T_b) := \operatorname{tr}[\hat{R}^{\dagger}(T_a)\hat{R}(T_b)] = (p^N)^{Q - |T_a, T_b|}$$
(26)

for any $T_a, T_b \in \Sigma_Q(p)$, where $\hat{R}(T)$ is defined in (25). The metric satisfies the usual properties $|T_a, T_b| \ge 0$, $|T_a, T_b| = |T_b, T_a|$, and $|T_a, T_b| = 0$ if and only if $T_a = T_b$. Additional details on the definition of this metric are provided in Appendix E.

The other property we need is that the metric in (26) is invariant¹¹ under the action of the direct product of two copies of the stochastic orthogonal group $\mathcal{O}_Q(p) \times \mathcal{O}_Q(p)$, acting as

$$|T_1, T_2| = |T_1', T_2'|, \text{ where } T_1, T_2 \in \Sigma_Q(p), (27)$$

and

$$T_1' = O_L T_1 O_R^{-1}, \quad T_2' = O_L T_2 O_R^{-1}, \quad \text{with } O_L, O_R \in \mathcal{O}_Q(p).$$
(28)

2. Stat mech model for the Clifford (stabilizer) random tensor network

The construction of the Clifford RTN follows the same steps as those reviewed in Sec. II A for Haar unitaries. We consider the RTN where on each edge there is an "on-site" Hilbert space $(\mathbb{C}^p)^M$ of dimension $d = p^M$ of M qudits, each qudit having a p-dimensional Hilbert space where p is a prime number. (For a brief review, see, e.g., Appendix B.) In complete analogy with Sec. II A, a Hilbert space $(\mathbb{C}^p)^{M \cdot z} = (\mathbb{C}^p)^N$ of dimension $D = p^N$, where N = Mz is associated with each vertex v, where z is the coordination number of a vertex.

Specifically, in the Clifford case the first line of (9) will be replaced by

$$\otimes^{\mathbb{Q}}(|T_v\rangle \langle T_v|) = \otimes^{\mathbb{Q}}[(\hat{V}|0\rangle^{\otimes N}) (\langle 0|^{\otimes N}\hat{V}^{\dagger})], \qquad (29)$$

where \hat{V} is an element of the Clifford group Cliff(N, p) acting on the Hilbert space (\mathbb{C}^p)^N at vertex v. This means simply that each¹² of the states $|1\rangle$ that appears in each of the Qtensor factors in (9) is replaced by the state $|0\rangle^{\otimes N} \in \mathbb{C}^N$, which is written in the computational basis, where $0 \in \mathbf{F}_p$ denotes the number zero in the finite number field \mathbf{F}_p . [See, e.g., the paragraph preceding that containing (23), or Appendix B.] The reason why $|0\rangle^{\otimes N}$ is chosen at each vertex is because this state is a stabilizer state.¹³ Owing to the fact that this state is a stabilizer state, the average of (29) over the Clifford group at a vertex v is (as shown in Ref. [65]) an equal-weight superposition over elements of the commutant [see, e.g., (B5) of Appendix B]:

$$\mathbb{E}_{\hat{V} \in \text{Cliff}(N,p)} [\otimes^{\mathcal{Q}} ((\hat{V}|0\rangle^{\otimes N})(\langle 0|^{\otimes N}\hat{V}^{\dagger}))]$$

= $\frac{1}{Z_{n,p,\mathcal{Q}}} \sum_{T_v \in \Sigma_{\mathcal{Q}}(p)} \hat{R}_v(T_v),$ (30)

where

$$\hat{R}_v(T) = [\hat{R}_e(T)]^{\otimes z}, \quad \text{with } \hat{R}_e(T) = [\hat{r}(T)]^{\otimes M}.$$

Then, in complete analogy with (20), one obtains using a maximally entangled state in an orthonormal basis on each

¹⁰Note that (as briefly reviewed below) in the stat mech models for the RTN and the quantum circuits monitored by measurements, Q is a replica (Rényi) index which goes to $Q \rightarrow 0$ in the RTN [56] and goes to $Q \rightarrow 1$ in the monitored quantum circuits [25]. Thus the inequality required for stabilization will be automatically satisfied in these limits of interest.

¹¹See (A11) of Appendix A for details.

¹²Recall that, in the Haar case, $|1\rangle$ is just one of the $D = d^z$ states of an orthonormal basis $|\mu_v\rangle$, $\mu_v = 1, ..., D$ that can occur at a vertex v.

¹³In fact any stabilizer state could be chosen at a vertex v; see Appendix B.

edge, a contribution to the partition function from that edge of the form

$$\frac{1}{(p^M)^Q} \operatorname{tr}[\hat{R}_e^{\dagger}(T_v)\hat{R}_e(T_{v'})] = (p^M)^{-|T_v, T_{v'}|},$$

where T_v and $T_{v'}$ denote elements of the commutant from the equal-weight sums (30) at the two vertices v and v' connected by the edge e. Then we obtain, by collecting all terms from the equal-weight sums appearing at each edge, in complete analogy to (22), the following (bulk) partition function of the Clifford RTN model:

$$Z_{\emptyset} = \mathbb{E}_{\hat{V} \in \text{Cliff}(M,p)} \text{tr} \rho^{\otimes Q}$$
(31)

$$= \prod_{\{T_v \in \Sigma_Q(p)\}} \exp\left\{-\sum_{\langle v, v' \rangle} \mathcal{H}(T_v, T_{v'})\right\},$$
$$\mathcal{H}(T_v, T_{v'}) = \ln(p^M)|T_v, T_{v'}|.$$
(32)

Observables, including the entanglement entropy, are formulated in the same way as in the Haar case. [See (5)–(7), and the discussion below (22).]

We recall from Ref. [56]¹⁴ that the transition in the RTN model can be driven by making (at each edge) the bond dimension random. In the Clifford case, this means that the bond dimension is $D_{\text{bond}} = p^M$ with p = prime = fixed, where M is independently distributed at each edge according to some probability distribution. A simple choice for this would be random dilution, corresponding to a binary probability distribution for M, taking on two values M = 0 and $M = M_0 =$ fixed at random. This would drive a transition in the RTN model corresponding to the (same) universality class possessing (see below) symmetry group $\mathcal{O}_Q(p) \times \mathcal{O}_Q(p)$ for any choice of a (large enough) value M_0 .

We stress that owing to the invariance property (27) of the metric, the Clifford stat mech model is invariant under the $\mathcal{O}_O(p) \times \mathcal{O}_O(p)$ symmetry group —up to an additional \mathbb{Z}_2 semidirect factor that is also present in the Haar case and will be irrelevant to our purposes, see Appendix A. Note that this is a *different* symmetry group¹⁵ for different primes p, but that this symmetry group is the same for all on-site Hilbertspace dimensions $d = p^M$ for a *fixed* prime *p*, independent of M: the symmetry group depends only on the prime p. This thus implies that any universal quantities occurring at continuous phase transitions in this stat mech model will in general depend on the prime p but will be the same for all on-site Hilbert-space dimensions $d = p^M$ of the same prime number p, independent of M. Note that this is in contrast with the Haar case, where the same universality class occurs for the transition at any finite on-site Hilbert-space dimension. In the Haar case, this must of course be the case, because universal properties cannot depend on short-distance ("ultraviolet") physics-all those stat mech models have the same



FIG. 2. The architecture of the monitored quantum circuit model. Each blue block represents a random unitary gate, and each hollowed circle represents a rank-1 projective measurement, occurring randomly with probability p_0 . The dynamical state $\rho(t)$ at circuit depth *t* is supported on qudits at the final time step (highlighted with purple squares) and is labeled by the measurement record **m**, see Eq. (33). We consider two cases, depending on whether the random unitaries are sampled from the Haar measure on U(D = 2d) (the "random Haar circuit"), or from the uniform distribution the Clifford group Cliff(2M, p) $\subset U(D)$ (the "random Clifford circuit").

symmetry.¹⁶ While this feature is the same in the Clifford case for a *fixed* value of the prime p (i.e., for on-site Hilbert-space dimension $d = p^M$ and different values of M), RTN models possessing on-site Hilbert-space dimensions which are powers of *different* primes have in general different universal critical properties at continuous phase transitions because they are invariant under different symmetry groups.

III. MEASUREMENT-INDUCED PHASE TRANSITION IN CLIFFORD CIRCUITS

A. Setup and replica trick

We now turn to measurement-induced entanglement transitions in monitored quantum circuits. We consider a one-dimensional chain of qudits with on-site Hilbert-space dimension d, subject to discrete-time dynamics generated by a quantum circuit with a "brick-wall" pattern, as illustrated in Fig. 2. Each unitary gate is acting on a pair of neighboring qudits and will be drawn from the Clifford group (we also briefly review the Haar case below). After each layer of the circuit (single time step), every site is either measured (projectively) with probability p_0 , or left untouched with probability $1 - p_0$. For a fixed set of unitary gates and measurement locations, the hybrid nonunitary dynamics of the system is described through the quantum channel

$$\rho(t) = \sum_{\mathbf{m}} K_{\mathbf{m}} \rho K_{\mathbf{m}}^{\dagger}, \qquad (33)$$

where ρ is the system's density matrix, **m** denotes measurement outcomes, and $K_{\rm m}$ are Kraus operators consisting of the random unitary gates and the projectors onto the measurement outcomes. We are interested in the entanglement properties of individual quantum trajectories $\rho_{\rm m} = K_{\rm m}\rho K_{\rm m}^{\dagger}$, which occur with the Born probability $p_{\rm m} = \text{tr}\rho_{\rm m}$. Our goal is to compute

¹⁴See Sec. IV F of Ref. [56].

¹⁵Recall that $\mathcal{O}_Q(p)$ denotes orthogonal matrices whose entries are elements of the number field \mathbf{F}_p , which itself has only pelements. Thus, the groups $\mathcal{O}_Q(p)$ are different finite groups for different primes p—they even have different order (=number of group elements).

¹⁶Which we recall from Ref. [56] is $S_Q \times S_Q$.

the Rényi entropy of such single trajectories, averaged over measurement outcomes and random unitary gates

$$S_A^{(n)} = \mathbb{E}_U \sum_{\mathbf{m}} p_{\mathbf{m}} \frac{1}{1-n} \ln \frac{\operatorname{tr} \rho_{A,\mathbf{m}}^n}{(\operatorname{tr} \rho_{\mathbf{m}})^n}.$$
 (34)

Ultimately, we also want to average over measurement locations and denote $\overline{S_A^{(n)}}$ the Rényi entropy averaged over measurement locations as well. As in the random tensor networks section above, we follow Refs. [10,24,25,56] and use a replica trick to compute (34):

$$\overline{S_A^{(n)}} = \mathbb{E}_U \sum_{\mathbf{m}} \lim_{k \to 0} \frac{\mathrm{tr}\rho_{\mathbf{m}}}{(1-n)k} \Big[\big(\mathrm{tr}\rho_{A,\mathbf{m}}^n\big)^k - \big(\mathrm{tr}\rho_{\mathbf{m}}^{\otimes kn}\big) \Big].$$
(35)

We write Q = nk + 1 the number of replicas, where the additional replica compared with the random tensor networks case comes from the Born probability weighting different quantum trajectories.

B. Generalized Weingarten functions

1. Haar case

Let us first briefly summarize the main ingredient of the statistical mechanics model of the measurement-induced transition, with Haar random unitary gates acting on qudits. Upon replicating the model, averaging over Haar gates drawn from the unitary group $U(D = d^2)$ naturally leads to degrees of freedom living in the permutation group S_Q , with Q = nk + 1the number of replicas. This follows from the Schur-Weyl duality detailed in the previous section, which states that the permutation group S_Q and the unitary group U(D) act on $(\mathbf{C}^D)^{\otimes Q}$ as a commuting pair. The Haar average of the replicated unitaries is given by (see Appendix C)

$$\mathbb{E}_{U \in U(D)}(U^{*Q} \otimes U^{Q}) = \sum_{\sigma, \tau \in S_{Q}} Wg_{D}(\sigma^{-1}\tau)\hat{R}_{v}(\sigma) \otimes \hat{R}_{v}(\tau),$$
(36)

where Wg are Weingarten functions, $D = d^2$, $\hat{R}_v(\sigma) = \hat{R}_e(\sigma) \otimes \hat{R}_e(\sigma)$ [see Eq. (12)] permutes the output legs of U by σ and contracts them with the corresponding legs of U^* [and similarly for $\hat{R}_v(\tau)$ acting on the input legs]. The tensor product in the right-hand side of Eq. (36) is over output and input legs of the unitary. Now contrary to the previous section on random tensor networks, the Weingarten function will directly enter the Boltzmann weights of the statistical models. As we review in Appendix C, those functions satisfy

$$\sum_{\tau \in S_Q} Wg_D(\sigma^{-1}\tau) D^{C(\tau)} = \delta_{\sigma,1},$$
(37)

where we recall that $C(\tau)$ denotes the number of cycles in the permutation τ . This implies that the Weingarten function is the inverse of the cycle counting function $D^{C(\tau)}$. Equation (36) naturally defines the degrees of freedom of the statistical model living on a honeycomb lattice (Fig. 3), where permutations live on vertices. Contracting unitary gates can be done by assigning a weight to links connecting unitaries given by

$$W(\sigma,\tau) = \operatorname{tr} \left[\hat{R}_{e}^{\dagger}(\sigma)\hat{R}_{e}(\tau)\right] = d^{C(\sigma^{-1}\tau)}.$$
(38)

Note the factor of d here, instead of D, since we are focusing on a single leg of the unitary. This weight is associated with all



FIG. 3. Statistical mechanics model for monitored quantum circuits. (a) The average over replicated unitary gates leads to degrees of freedom belonging to the commutant, defined on vertices; Weingarten weights Wg_D , represented by the dashed line; and link weights W, corresponding to purple solid lines. (b) The statistical mechanics model is defined on an anisotropic honeycomb lattice.

links that were not measured. If a link was measured instead, all replicas are constrained to be in the same state, and the weight is instead d after averaging over possible measurement outcomes. Those equations fully determine the weights of the statistical model in monitored Haar random circuits. Upon averaging over measurement locations and outcomes, the weight assigned to a link is therefore given by [25]

$$W_p(\sigma, \tau) = (1 - p_0)d^{C(\sigma^{-1}\tau)} + p_0d.$$
 (39)

Putting these results together, we obtain an anisotropic statistical mechanics model defined on the honeycomb lattice,

$$Z = \sum_{\{g_i \in S_Q\}} \prod_{\langle ij \rangle \in G_s} W_p(\sigma_i^{-1}\tau_j) \prod_{\langle ij \rangle \in G_d} Wg_D(\sigma_i^{-1}\tau_j), \quad (40)$$

where G_s (G_d) denotes the set of solid (dashed) links on the lattice. In Fig. 3, the vertical (dashed) links on the honeycomb lattice represent the Weingarten functions which originated from averaging the two-site unitary gates, and the solid links keep track of the link weights originating from averaging over measurements.

2. Clifford case

Let us now turn to the Clifford case, with random unitary gates drawn from the Clifford group Cliff $(N, p) \subset U(D)$, with $D = d^2$, N = 2M, and $d = p^M$ with p prime. As in the Haar case, the average over Clifford gates is given in terms of tensor products of elements of the commutant of the Clifford group (see Appendix D)

. 0

$$\mathbb{E}_{U \in \operatorname{Cliff}(N,p)}(U^{*\mathcal{Q}} \otimes U^{\mathcal{Q}})$$

= $\sum_{T_i, T_o \in \Sigma_{\mathcal{Q}}(p)} \operatorname{Wg}_D(T_i, T_0) \hat{R}_v(T_o) \otimes \hat{R}_v(T_i),$ (41)

where, as before, $\hat{R}_v(T_i) = \hat{R}_e(T_i) \otimes \hat{R}_e(T_i)$ is acting on incoming legs, while $\hat{R}_e(T_o)$ acts on outgoing legs, where the operator $\hat{R}_e = \hat{r}^{\otimes M}$ was introduced in the previous section, see Eq. (23). Here the coefficients $Wg_D(T_i, T_0)$ are generalized Weingarten weights. The procedure to determine them follows closely the Haar case, and we show in Appendix D that the generalization of Eq. (37) reads

$$\sum_{T_b \in \Sigma_Q(p)} Wg_D(T_a, T_b) tr[\hat{R}_v^{\dagger}(T_b)\hat{R}_v(T_c)] = \delta_{T_a, T_c}, \qquad (42)$$

as well as $\sum_{T_b \in \Sigma_Q(p)} \operatorname{tr}[\hat{R}_v^{\dagger}(T_a)\hat{R}_v(T_b)] \operatorname{Wg}_D(T_b, T_c) = \delta_{T_a, T_c}$. In other words, the Weingarten weights (when viewed as a matrix) are given by the inverse of the matrix $\operatorname{tr}[\hat{R}_v^{\dagger}(T_a)\hat{R}_v(T_b)]$. On the other hand, the link weights connecting two vertices are given by

$$W(T_a, T_b) = \text{tr}[\hat{R}_e^{\dagger}(T_a)\hat{R}_e(T_b)] = d^{Q - |T_a, T_b|}, \qquad (43)$$

with $d = p^M$. As in the Haar case, this weight is replaced by a factor of d if a measurement occurred on that link (upon averaging over measurement outcomes), as the measurement constrains all replicas to be in the same state. After averaging over measurement locations, the weight assigned to a link reads

$$W_p(T_a, T_b) = (1 - p_0)d^{Q - |T_a, T_b|} + p_0 d.$$
(44)

The partition function of the statistical model in the Clifford case is then given by

$$Z = \sum_{\{T_i \in \Sigma_Q(p)\}} \prod_{\langle ij \rangle \in G_s} W_p(T_i, T_j) \prod_{\langle ij \rangle \in G_d} Wg_D(T_i, T_j), \quad (45)$$

with the same lattice as in the Haar case (depicted in Fig. 3).

Note that both the link and Weingarten weights are invariant under two copies of the stochastic orthogonal group $\mathcal{O}_{\mathcal{Q}}(p) \times \mathcal{O}_{\mathcal{Q}}(p)$, acting as $T'_{a} = O_{L}T_{a}O_{R}^{-1}$, $T'_{b} = O_{L}T_{b}O_{R}^{-1}$, with $O_{L}, O_{R} \in \mathcal{O}_{\mathcal{Q}}(p)$. The conclusions for the Clifford RTN statistical mechanics model thus carry over to Clifford monitored quantum circuits: the Clifford stat mech model is invariant under a global $\mathcal{O}_O(p) \times \mathcal{O}_O(p)$ symmetry group. This symmetry group depends only on the prime number p, and not on M in the on-site Hilbert-space dimension $d = p^{M}$. This implies that all universal quantities of the entanglement transition in Clifford monitored circuits will depend on p, but will be independent of M. This is in sharp contrast to the Haar case, where the same universality class occurs for the transition at any finite on-site Hilbert-space dimension. Although the statistical mechanics model (45) describing the monitored Clifford circuit has the same symmetry group as the one describing stabilizer RTNs (32), both problems correspond to different replica limits: $Q \rightarrow 0$ for RTNs, and $Q \rightarrow 1$ for monitored quantum circuits.

C. $M \rightarrow \infty$ limit with *p* fixed and percolation

While the Clifford statistical mechanics formulation is in general not analytically tractable, the limit of large on-site Hilbert space leads to drastic simplifications. Let us first consider the link weight, given by Eq. (43), where we recall that $|T_a, T_b|$ is a metric satisfying $|T_a, T_b| \ge 0$, $|T_a, T_b| = |T_b, T_a|$, and $|T_a, T_b| = 0$ if and only if $T_a = T_b$. This immediately

implies that, at large d, the weight for nonmeasured links simplifies dramatically to

$$W(T_a, T_b) \sim d^Q \delta_{T_a, T_b} + \cdots . \tag{46}$$

This corresponds to a perfectly ferromagnetic interactions that forces the elements of the commutant to be identical on unmeasured links. Since the Weingarten weights are given by the inverse of tr[$\hat{R}_v^{\dagger}(T_i)\hat{R}_v(T_o)$], we immediately find

$$Wg_D(T_i, T_0) \sim D^{-Q} \delta_{T_i, T_0} + \cdots, \qquad (47)$$

with $D = d^2$. Note that those results hold for $d = p^M \to \infty$, independently of how the limit is taken. We now focus on the limit of $d = p^M$ with p prime fixed and $M \to \infty$. (We discuss the limit of large p in the next section). Combined, those results tell us that in the $M \to \infty$ limit, the statistical mechanics weights for Clifford circuits become identical to those of the Haar case: precisely in the $M \to \infty$ limit we obtain a Potts model, whose states are the elements T_a of the commutant $\Sigma_Q(p)$. The number of states in the Potts model depends on p and on the number of replicas Q and is given by the dimension of the commutant [65]:

$$|\Sigma_{\mathcal{Q}}(p)| = \prod_{k=0}^{Q-2} (p^{k}+1) = p^{\frac{(Q-1)(Q-2)}{2}} \frac{\prod_{k=0}^{\infty} \left(1 + \frac{1}{p^{k}}\right)}{\prod_{k=0}^{\infty} \left(1 + \frac{1}{p^{k+Q-1}}\right)}.$$
(48)

(In the second equality, obtained by straightforward rewriting, the number Q can be analytically continued from integer values.) In the replica limit $Q \rightarrow 1$, we have $|\Sigma_Q(p)| \rightarrow 1$, so the critical theory of the measurement induced transition is simply described by classical percolation. In particular, this predicts a diverging correlation length with a critical exponent v = 4/3. Note that the way the replica limit is approached depends on p. This shows up in, e.g., the effective central charge $c_{\text{eff}} = \lim_{Q \rightarrow 1} \frac{1}{dc} \frac{1}{dQ}$ introduced in Ref. [49]. The central charge as a function of the number of replicas Q is now given by $c(Q) = 1 - \frac{6}{x(x+1)}$ with $x + 1 = \pi / \arccos(\sqrt{|\Sigma_Q(p)|}/2)$. This leads to a closed form expression for the effective central charge:

$$c_{\text{eff}}^{M \to \infty} = \frac{5\sqrt{3}}{8\pi} \bigg[2\psi_{\frac{1}{p}} \bigg(\frac{i\pi}{\ln p^{-1}} \bigg) - \ln \frac{p^3}{(p-1)^2} \bigg], \qquad (49)$$

where $\psi_q(z)$ is the *q* digamma function, which is defined as the derivative of $\ln\Gamma_q(z)$ with respect to *z*, where $\Gamma_q(z)$ is the *q*-deformed Gamma function. The special case p = 2 of this formula was reported in Ref. [49].

This mapping to percolation is specific to the $M \to \infty$ limit: if M is large but finite, the critical theory of the measurement-induced transition is described by the percolation CFT perturbed by a relevant perturbation (identified as the "two-hull" operator in Refs. [25,56]), with scaling dimension $\Delta = 5/4$. To see this, let us consider the Landau-Ginzburg formulation of the $|\Sigma_Q(p)|$ -state Potts field theory in terms of the Potts order parameter field ϕ_a where $a = 1, \ldots, |\Sigma_Q(p)|$ labels elements of the commutants, and $\sum_a \phi_a = 0$. The symmetry of the Potts theory for $M \to \infty$ is $S_{|\Sigma_Q(p)|}$, which is much larger than the stochastic orthogonal symmetry $\mathcal{O}_Q(p) \times \mathcal{O}_Q(p)$ of the finite-M case. To see this, note that $\mathcal{O}_Q(p) \times \mathcal{O}_Q(p) \subset S_{|\mathcal{O}_Q(p)|}$ using Cayley's theorem: the left and right actions of the group $\mathcal{O}_Q(p)$ on itself have a permutation representation. Note also that $S_{|\mathcal{O}_Q(p)|}$ is a subgroup of $S_{|\Sigma_Q(p)|}$ since the stochastic orthogonal group is a subset of the commutant $\Sigma_Q(p)$. The symmetry-breaking $S_{|\Sigma_Q(p)|} \rightarrow \mathcal{O}_Q(p) \times \mathcal{O}_Q(p)$ can be implemented by the perturbing the Potts model,

$$\mathcal{L} = \mathcal{L}_{\text{Potts}} + \sum_{g,h \in \mathcal{O}_Q(p)} V(g^{-1}h)\phi_g\phi_h,$$
(50)

where V is a class function of the group $\mathcal{O}_Q(p)$, ensuring a left and right stochastic orthogonal group symmetry. The perturbation has scaling dimension $\Delta = 5/4$ at the Potts (percolation) fixed point in the replica limit and is therefore relevant.

The fate of this theory in the IR is not known analytically and corresponds to the generic universality class of Clifford circuits for a given p. For p fixed and M large but finite, we thus expect a crossover between percolation and the finite $d = p^M$ universality class (dependent on p), with the corresponding crossover length scale

$$\xi(M) \sim p^{4M/3},$$
 (51)

where we have assumed that the relevant perturbation comes with an amplitude $V \sim d^{-1}$. Note that this crossover length scales exponentially with M, giving a very broad regime $L \ll$ $\xi(M)$ described by percolation at large M. For length scales much smaller than $\xi(M)$, we expect to see percolation physics, and in particular, the entanglement entropy should be given by a minimal cut picture as we derive in the next section. In particular, the entanglement entropy in that regime scales with $\ln d = M \ln p$ (see next section). However, for length scales much larger than $\xi(M)$, the physics is controlled by the infrared fixed point of the theory (50), corresponding to the generic Clifford universality class for fixed prime p. In particular, in this regime, the prefactor of the logarithmic scaling of the entanglement entropy at criticality will be universal, and will depend on p, but not on M.

D. Large-*p* limit and minimal cut

Finally, we now comment on the large-p (prime) limit, keeping M = 1 fixed. For each finite p, there is a distinct universality class, which is independent of M. However, the results of the previous section also indicate that the bulk theory should approach percolation as $p \to \infty$ (with M = 1fixed): we thus expect a series of fixed points approaching percolation as $p \to \infty$. A simple way to understand this is to consider a fixed realization of measurement locations. Each bond that is measured is effectively cut, while all other weights constrain the statistical model's spins to be the same in this limit. We thus obtain a simple percolation picture of fully ordered (zero temperature) ferromagnetic spin model diluted by the measurements. As we show below, a frustrated link costs a large energy $\approx \ln p$, leading to an effective minimal cut picture in that limit.

Recall that computing entanglement requires computing two different partition functions Z_A and Z_{\emptyset} , which differ only by their boundary condition on the top boundary of the circuit. The boundary condition for the calculation of Z_A forces a different boundary condition in region A, and thus introduces a domain wall (DW) near the top boundary. In the limit $d \to \infty$, the DW is forced to follow a minimal cut, defined as a path cutting a minimum number of unmeasured links (assumed to be unique for simplicity): the argument follows closely Ref. [50] in the Haar case. Due to the uniform boundary condition in Z_{\emptyset} , all vertex elements in Z_{\emptyset} are equal, so Z_{\emptyset} is trivial and give by a single configuration of spins. Z_A differs from Z_{\emptyset} due to the fact the DW will lead to frustrated links that contribute different weights to Z_A . Each frustrated unmeasured link contributes a factor tr $[\hat{r}^{\dagger}(1)\hat{r}(g_{SWAP})]/\text{tr} \hat{r}(1)$ to the ratio of partition functions Z_A/Z_{\emptyset} , where $g_{SWAP} = (1, 2, ..., n)^{\otimes k}$ is the boundary condition enforced in the entanglement region *A* in Z_A . (The permutation g_{SWAP} acts trivially on the last replica implementing the Born probability factor.) We find tr $[\hat{r}^{\dagger}(1)\hat{r}(g_{SWAP})]/\text{tr} \hat{r}(1) = p^{k+1-Q} = p^{(1-n)k} \ll 1$. This corresponds (since we are considering large *p*) to a very large energy cost per frustrated link

$$\Delta E = (n-1)k\ln p,\tag{52}$$

so the domain wall will follow a path through the circuit minimizing the number of unmeasured links it has to cut. This leads to the expression $Z_A = p^{(1-n)k \cdot \ell_{\text{DW}}} Z_{\emptyset}$, with ℓ_{DW} the number of unmeasured links that the DW crosses along the minimal cut. In the replica limit, this leads to a simple expression for the Rényi entropies:

$$S_A^{(n)} = \ell_{\rm DW} \ln p, \tag{53}$$

where this equation is valid for any given configuration of measurement locations. We use $\overline{\ell_{DW}}$ to denote the average of ℓ_{DW} over measurement locations, which are simply percolation configurations. This quantity has a simple scaling in percolation: it is extensive $\overline{\ell_{DW}} \sim L_A$ (volume law) for $p_0 < p_{0,c} = 1/2$, and constant $\overline{\ell_{DW}} \approx O(1)$ (area law) for $p_0 > p_{0,c} = 1/2$. At criticality, this implies a logarithmic scaling of the entanglement entropy [14,68,69]

$$\overline{S_A^{(n)}} \underset{p\gg1}{\approx} \frac{\sqrt{3}}{\pi} \ln p \ln L_A.$$
(54)

We expect the measurement-induced transition to approach these predictions at large p.

E. Forced measurements and random tensor networks at large bond dimension

Our predictions can also be generalized to the case of forced measurements, where the given outcome for a "forced measurement" is chosen randomly in a way that is independent of the state, instead of following the Born rule. In the context of RTNs, entanglement transitions can be implemented at fixed bond dimension by randomly breaking bulk bonds [56], which can be thought of as such "forced measurements" [34]. The corresponding statistical mechanics models have the same stochastic orthogonal group symmetry but correspond to different replica limits $Q \rightarrow 1$ (projective measurements) vs $Q \rightarrow 0$ (forced measurements, RTNs), and are thus in general expected to be in different universality classes. However, numerical results suggest that forced vs projective measurements might be in the same universality class in the Clifford case [58], indicating that the two replica limits $Q \rightarrow 0$ and $Q \rightarrow 1$ could lead to the same critical theory.

The conclusions of the previous sections carry over to this forced measurement setup as well: in the large-D limit,

Data at the critical point	d = 2	d = 3	d = 5	<i>d</i> = 97	d = 997
$p_{0,c}$	0.160	0.278	0.377	0.495	$0.500 \approx 1/2$
$h_{a b}$	0.53	0.62	0.66	$1.26 \approx \frac{\sqrt{3}}{2\pi} \ln(97)$	$1.90 \approx \frac{\sqrt{3}}{2\pi} \ln(997)$
h _{MN}	3.0	2.9	2.6	2.2	2.0
$h_{f f}^{(1)}$	0.41				$0.33 \approx 1/3$

TABLE I. A comparison of the location of the critical point $p_{0,c}$ and the operator scaling dimensions $h_{a|b}$, h_{MN} , $h_{f|f}^{(1)}$ in hybrid Clifford circuits with different prime qudit dimensions d = p.

the link and Weingarten weights simplify dramatically, the statistical mechanics model reduces to a Potts model, and the large d = p limit also satisfies the minimal cut prediction (54). The only difference is that the number of replicas is Q = nk instead of Q = nk + 1, reflecting the absence of Born probabilities. In particular, the replica limit corresponds to $Q \rightarrow 0$ instead of $Q \rightarrow 1$. At large p and Q small (i.e., 0 < Q < 1), the dimension of the commutant reads, using Eq. (48),

$$|\Sigma_{\varrho}(p)| \underset{p \to \infty}{\sim} \frac{2}{1+p^{\varrho}} p^{\frac{\varrho(\varrho+1)}{2}}, \tag{55}$$

which gives $|\Sigma_Q(p)| \to 1$ in the replica limit $Q \to 0$. At large *p*, the replica limit is thus given by a Potts model with $|\Sigma_Q(p)| \to 1$ states, consistent with percolation and the minimal cut prediction.

IV. NUMERICAL RESULTS

In this section we provide numerical evidence that supports our key conclusions from the previous sections on symmetry, namely, that the universality class of the transition for qudit dimension $d = p^M$ depends on the prime number p, but not on the power M. To this end, we consider two cases:

(A) $d = p^M$ with fixed M = 1 and different values of p, where we expect a series of different, p-dependent universality classes;

(B) $d = p^M$ with fixed p and a few small values of M, where we expect to obtain the same exponents while varying M.

For each value of d, we simulate monitored Clifford circuits of L = 512 qudits with periodic boundary condition and focus on the steady-state entanglement properties at their respective critical points. The circuit thus takes the geometry of a semi-infinite cylinder, and at criticality the entanglement entropy of a region A is predicted to take the following form:

$$S(A = [z_1, z_2]) = 2h_{a|b} \ln\left(\frac{L}{\pi} \sin\frac{\pi z_{12}}{L}\right) \coloneqq 2h_{a|b} \ln w_{12},$$
(56)

where $h_{a|b}$ is the scaling dimension of a (primary) boundarycondition changing operator in the underlying CFT.

A. Different universality classes at various $d = p^{M=1}$

Here, we take the qudit dimension to be a prime number, $d = p^{M=1}$ for $p \in \{2, 3, 5, 97, 997\}$. In Fig. 4(a), we plot the results of $S(A = [z_1, z_2])$ in the form of Eq. (56), and extract the scaling dimension $h_{a|b}$ at the respective critical points. The locations of the critical points are determined by the best fit of $S(A = [z_1, z_2])$ to a linear function of $\ln w_{12}$, see Eq. (56). The fits for $h_{a|b}$ and $p_{0,c}$ are collected in Table I.

From our fitting, the scaling dimension $h_{a|b}$ clearly depends on p, and increases monotonically with p. It approaches the prediction from minimal cut percolation at large p, namely, $h_{a|b} = \frac{\sqrt{3}}{2\pi} \ln p$ —see Eqs. (54) and (56). In addition, the value of $p_{0,c}$ approaches 1/2 as p is increased, again consistent with bond percolation on a square lattice.¹⁷

We also include results of the "mutual negativity" N(A, B)between two disjoint regions $A = [z_1, z_2]$ and $B = [z_3, z_4]$ [70]. This quantity is expected to take the following form when the cross ratio η for the four endpoints is small,

$$N(A = [z_1, z_2], B = [z_3, z_4]) \propto \eta^{h_{\rm MN}}, \text{ as } \eta \to 0.$$
 (57)

Here, $h_{\rm MN}$ stands for the "mutual negativity exponent." In Fig. 4(b) and Table I, we see that $h_{\rm MN}$ is close to three for smaller values of d = p, and approaches two as d = p becomes large, as consistent with minimal cut percolation [70].

To further contrast universality classes at small d = p with percolation at large d = p, we focus on the maximum and minimum qudit dimensions in our numerics, namely, d = $p \in \{2, 997\}$, and consider yet another quantity, known as a localizable entanglement [71], recently introduced in the context of hybrid circuits [21]. In this case, we again choose two disjoint regions $A = [z_1, z_2]$ and $B = [z_3, z_4]$ and perform a single-qudit projective measurement on each qudit outside $A \cup B$. In the postmeasurement state, the mutual information between A and B is expected to be a conformal four-point function [21], and takes the following form when the cross ratio η for the four endpoints is small:

$$I(A = [z_1, z_2], B = [z_3, z_4]) \propto \eta^{h_{f/f}^{(1)}}$$
 as $\eta \to 0.$ (58)

Here, $h_{f|f}^{(1)}$ is another critical exponent, which takes the value of 1/3 in the percolation limit $d = p \rightarrow \infty$. In our numerics [see Fig. 4(c)], we find that $h_{f|f}^{(1)}$ fits well to $1/3 \approx 0.33$ at d = p = 997 but takes a *distinct* value ≈ 0.41 at d = p = 2 [21].

B. Same universality class for d = p and $d = p^2$

Here we compare qudit dimensions $d = p^M$ at a fixed p but different M. To probe the infrared fixed point, we choose M = 1, 2 so that $L \gg \xi(M)$, where $\xi(M)$ is a crossover length scale

¹⁷We note in passing that the series of $p_{0,c}$ as summarized in Table I violates a conjectured bound [18] for all values of $d = p \ge 3$ that we accessed.



FIG. 4. Numerical results for $d = p^{M=1}$. The fits for the critical exponents are summarized in Table I. (a) We plot $S(A = [z_1, z_2])$ against $\ln w_{12}$ [see Eq. (56)] at different values of d = p, where we see a clear dependence of the slope $(=2h_{a|b})$ on p. With increasing p, the value of $h_{a|b}$ approaches that of minimal cut percolation [see Eq. (54)]. (b) The mutual negativity N(A, B) between two disjoint subregions $A = [z_1, z_2]$ and $B = [z_3, z_4]$ against the cross ratio $\eta = (w_{12}w_{34})/(w_{13}w_{24})$. The power law at small η defines a critical exponent h_{MN} , which has a clear dependence on p, and approaches minimal cut percolation at large p. (c) A "localizable entanglement" between A and B. We perform a single-qudit projective measurements on each qudit outside $A \cup B$, and calculate the mutual information I(A, B) subsequently. The power law at small η is another critical exponent $h_{f|f}^{(1)}$, which should in general take different values at different p, as we illustrate.

that diverges with *M* [see Eq. (51)]. Numerically accessing the crossover from $L \gg \xi(M)$ to $L \ll \xi(M)$ is an interesting problem left for future versions of this work.



FIG. 5. Numerical results for $d = p^M$, where we take $p \in \{2, 3\}$ and $M \in \{1, 2\}$. Comparing Fig. 4(a) and Eq. (56), we see that $h_{a|b}$ depends on the choice of p but not on M.

For concreteness, we briefly describe the circuit model for M > 1. At each site of the lattice we construct a "composite" qudit of dimension $d = p^M$ by grouping together M "elementary" qudits each of dimension p. The circuit takes a brickwork architecture in the composite qudits, where nearest-neighbor composite qudits are coupled with random unitaries from the group Cliff(2M, p), and single composite qudits are subject to rank-1 projective measurements with probability p_0 .

We compute the entanglement entropy S(A) of region A for $d \in \{2, 4 = 2^2\}$ and for $d \in \{3, 9 = 3^2\}$. The results are plotted in Fig. 5, where we see that $h_{a|b}$ depends p but not on the choice of M. Results for $h_{a|b}$ at other values of p also follow this pattern but are not displayed.

Critical exponents for Clifford RTNs from Ref. [58] are summarized in Table II for comparison. Our numerical results indicate that the critical properties do not depend on the measurement scheme, forced (corresponding to the replica limit $Q \rightarrow 0$ in the statistical mechanics model) vs projective (replica limit $Q \rightarrow 1$) [58] but appear to differ (slightly) between monitored Clifford circuits and Clifford RTNs for small p. Since the statistical mechanics models for RTNs (31) and monitored circuits (45) have the same symmetry group, we expect the entanglement transitions in both cases to be in the same universality class for a given measurement scheme (forced or projective). The numerical discrepancy between those cases might be due to strong corrections to scaling (i.e., from irrelevant operators) in the RTN case for small p, since there is no transition for p = 2 in that case. One can, in principle, get an idea of the strength of such corrections by comparing Clifford RTNs with bond dimensions $d = p^{M}$ at a few small values of *M*, in a similar fashion to Fig. 5, although we leave a more thorough analysis of those corrections for future work.

C. Multifractal scaling of the purity

In this section we take the prime number to be p = 2. Based on the discussion and the results in the rest of this paper, we have no reason to expect that the general physics discussed in this section (multifractal scaling) will be fundamentally

Data at the critical point	d = 2	d = 3	d = 5	<i>d</i> = 23	d = 503
$h_{a b}$		$0.53 \approx 0.48 \ln(3)$	$0.61 \approx 0.38 \ln(5)$	$0.91 \approx 0.29 \ln(23)$	$1.74 \approx 0.28 \ln(503)$
h _{MN}		2.8	2.5	2.1	2.1
$h_{f f}^{(1)}$		0.37	0.36	$0.33 \approx 1/3$	$0.33 \approx 1/3$

TABLE II. Exponents for Clifford RTN [58] with different prime qudit dimensions d = p. The exponents are all close to the Clifford circuit but different.

different for other prime numbers p, except that numerical values for various critical exponents defined below are expected to depend explicitly on the prime p for the symmetry reasons mentioned in the other parts of this paper.

Specifically, in this section, we consider the purity of the normalized reduced density matrix [compare (34)]

$$\hat{\rho}_{A,\mathbf{m}} = \frac{\rho_{A,\mathbf{m}}}{\mathrm{tr}(\rho_{A,\mathbf{m}})}$$

of an interval *A* at the final time-slice of the circuit. In every realization of circuit disorder (arising from the quantum trajectory **m**, the Clifford unitaries and measurement locations, the dependence on which is completely or partially suppressed here and from now on, for ease of notation), the spectrum of $\hat{\rho}_{A,\mathbf{m}}$ is known to be flat:¹⁸ Thus, in every realization of disorder, the *n*th Rényi entropies $S_A^{(n)}$,

$$\operatorname{tr}(\hat{\rho}_{A,\mathbf{m}})^n = \exp\left\{-(n-1)S_A^{(n)}\right\},\$$

are all equal, and we consider without loss of generality the case n = 2. [Again, the dependence of $S_A^{(n)}$ on the quantum trajectory **m** and the other types of disorder is suppressed, which is also done for $G(x_1, x_2)$ defined below.]

Specifically, we consider the disorder averages,¹⁹ denoted by an overbar $\overline{(\ldots)}$, of powers (moments) of the purity

$$G(x_1, x_2) := \operatorname{tr}(\hat{\rho}_{A,\mathbf{m}})^2 = \exp\left\{-S_A^{(2)}\right\},$$
(59)

where x_1 and x_2 are the endpoints of the interval A, i.e., $|x_1 - x_2| = |A|$: At the transition, these moments will all scale with certain critical exponents X_k :

$$\overline{\left[G_{(x_1,x_2)}\right]^k} \sim \frac{B_k}{R_{12}^{2X_k}},\tag{60}$$

where R_{12} is the chord distance

$$R_{12} := \frac{L}{\pi} \sin\left(\frac{\pi}{L}|x_1 - x_2|\right),$$

L is the spatial size (with periodic boundary conditions), and B_k are nonuniversal amplitudes which turn out to be of little interest to us.²⁰ The moment averages in (60) are those of the two-point correlation function $G(x_1, x_2)$ associated with the so-called boundary-condition changing (bcc) operator twisting the boundary conditions [25,56] at the endpoints of the

interval *A*. If the dependence of the exponents X_k on the index *k* is *nonlinear*, the correlation function $G(x_1, x_2)$ is said to obey *multifractal* scaling [49,72,73]. (See Appendix F for a brief review.) This means in particular that the *mean* and the *typical* values of $G(x_1, x_2)$ scale with *different* critical exponents, and that the entanglement transition possesses a *continuous spectrum of critical exponents* [corresponding to X_k with continuous real values of *k*, arising from noninteger moment powers in (60)]. Whether multifractal scaling is present can be inferred from the scaling of the statistical fluctuations of $S_A^{(2)}$ as follows:

We know from (59) that, in every realization of disorder, the negative of the logarithm of $G(x_1, x_2)$ is the second Rényi entropy,

$$-\ln G(x_1, x_2) = S_A^{(2)} (|x_1 - x_2| = |A|).$$

As briefly reviewed in Appendix F, a comparison of the cumulant expansion of the left-hand side of Eq. (60) with the Taylor expansion of the exponents X_k in powers of k [appearing on the right-hand side of Eq. (60)] shows that all cumulants of the second Rényi entropy $S_A^{(2)}$ grow linearly with the logarithm of the chord distance, $\ln R_{12}$. The coefficients of proportionality are universal and equal to $(-1)^{k-1}$ times twice the Taylor coefficients of X_k in powers of k,

$$\overline{S_A^{(2)}} \sim 2x^{(1)} \ln R_{12},$$
 (61)

$$\left(S_A^{(2)} - \overline{S_A^{(2)}}\right)^2 \sim -2x^{(2)} \ln R_{12},$$
 (62)

$$\kappa_3[S_A^{(2)}] \sim 2 x^{(3)} \ln R_{12},$$

:, (63)

where $\kappa_3[S_A^{(2)}]$ denotes the third cumulant of the random variable $S_A^{(2)}$, and

$$X_k = k x^{(1)} + \frac{k^2}{2!} x^{(2)} + \frac{k^3}{3!} x^{(3)} + \cdots .$$
 (64)

Note that (61) is nothing but (35) expressed in terms of $G(x_1, x_2)$ defined in (59).

As we summarize below, we find numerically that the universal numbers $x^{(2)}$ and $x^{(3)}$, characterizing statistical fluctuations of the entanglement entropy about its mean, are both nonvanishing, and that at least $x^{(2)}$ is of a magnitude which is a sizable fraction of $x^{(1)}$, the latter characterizing the *averaged* entanglement entropy (35). It then follows from (64) that the dependence of the exponent X_k on the moment-order k is non-linear. As mentioned above, this signifies multifractal scaling of the random variable $G(x_1, x_2)$ defined in (59), i.e., it implies (in particular) different scaling exponents for the average and typical values of this quantity: The typical exponent describes

¹⁸All nonvanishing eigenvalues of $\hat{\rho}_{A,\mathbf{m}}$ are equal.

¹⁹Again, the average is over quantum trajectories involving the Born rule probability, over random unitaries and measurement locations.

²⁰The relationship of the nonuniversal amplitudes B_k with the universal results that we discuss below is mentioned in a footnote of Appendix F.



FIG. 6. Numerical results for the first three cumulants of the second Rényi entanglement entropy $S_A^{(2)} = S_{A=|x_1,x_2|}$: Averaged entanglement entropy (blue) from (61), its second cumulant (red) from (62), and its third cumulant (green) from (63), versus the chord distance, here denoted by R_{12} . The slopes of the plots yield the exponent $x^{(1)} = h_{a|b} \approx 0.53$, $-x^{(2)} = h_{a|b}^{(2)} \approx 0.32$, and $x^{(3)} = h_{a|b}^{(3)} \approx 0.15$. (The notations $h_{a|b}$, $h_{a|b}^{(2)}$, and $h_{a|b}^{(3)}$ serve as a reminder that these are the exponents associated with the boundary-condition changing (bcc) operator [21,25,56].)

the universal prefactor factor $x^{(1)}$ multiplying the logarithm of subsystem size of the *averaged entanglement entropy*, as in (61). The exponent $X_1 \ (\neq x^{(1)})$ describes the algebraic decay of the *averaged purity* [from (59) and (60) with k = 1],

$$\overline{\operatorname{tr}(\hat{\rho}_{A,\mathbf{m}})^2} = \overline{G(x_1, x_2)} \sim \frac{B_1}{R_{12}^{2X_1}}$$

We now proceed to discuss the numerical results. In Fig. 6 we plot the numerical values of the first three cumulants of the entanglement entropy $S_A^{(2)} = S_{A=|x_1-x_2|}$ versus the chord distance R_{12} in the figure: From the slopes of the plots of the averaged entanglement entropy (blue) from (61), of its second cumulant (red) from (62), and of its third cumulant (green) from (63), we read off the critical exponents $x^{(1)} = h_{a|b} \approx 0.53$, and $-x^{(2)} = h_{a|b}^{(2)} \approx 0.32$, and $x^{(3)} = h_{a|b}^{(3)} \approx 0.15$, respectively. (These notations $h_{a|b}, h_{a|b}^{(2)}$, as well as $h_{a|b}^{(3)}$, just serve as a reminder that these are the exponents associated with the boundary-condition changing (bcc) operator [21,25,56], using the notation of Ref. [21].)

Given the numerical values of these exponents we see that the expansion term of X_k in (64) quadratic in k is of a sizable fraction [$\approx 0.32/(2 \times 0.53) = 0.302$] of the term linear in k. Thus we conclude that X_k is clearly not a linear function of kand that the purity thus exhibits multifractal scaling. The expansion term of X_k cubic in k, even though small as compared with $x^{(1)} = h_{a|b}$, does not appear entirely negligible, which is presumably indicative of a k dependence of X_k beyond quadratic order.

In conclusion, our numerical study of the statistical fluctuations of the entanglement entropy has established beyond doubt multifractal scaling of the purity in our Clifford circuit for prime p = 2. Moreover, as mentioned at the beginning of this section, we have no reason to doubt that the scaling behavior of the purity is also multifractal for on-site Hilbertspace dimensions $d = p^M$ with other prime numbers p, except that also the exponents $x^{(2)}, x^{(3)}, \ldots$, in addition to $x^{(1)} = h_{a|b}$ (see Table I), are expected to depend explicitly on the prime p, based on our results in the rest of this paper.

V. DISCUSSION

In this work, we have introduced a replica statistical mechanics mapping to compute entanglement properties of stabilizer random tensor networks, and monitored Clifford quantum circuits acting on qudits with $d = p^{M}$. The degrees of freedom (spins) of the resulting statistical models belong to the commutant of the Clifford group, using a generalization of the concept of Schur-Weyl duality to the Clifford case. We find that the Boltzmann weights are invariant under the left and right action of the stochastic orthogonal group with entries in the finite number field \mathbf{F}_p . This implies, surprisingly, that the universal properties of the entanglement transitions in Clifford circuits and RTNs depend on the on-site Hilbert-space dimension d but only through the prime p. This is in sharp contrast with the Haar case where the symmetry group is independent of d as long as it is not infinite. Our formalism also allowed us to derive exact mappings onto classical percolation and minimal cut formulas as $d = p^M$ becomes large, with different consequences for p fixed and $M \to \infty$, or M fixed and $p \to \infty$. We confirmed our predictions using large-scale stabilizer numerics, which also showed that projective and forced measurement schemes yield the same finite-d critical exponents for Clifford circuits.

Our large-*d* results here bolster a known result about stabilizer tensors. Random stabilizer tensors with large bond dimension are perfect tensors with high probability [59,74,75]. A perfect tensor has the special property of being a unitary gate for any equal-size bipartition of its legs. Thus, a Clifford circuit or RTN as $d \rightarrow \infty$ becomes an assembly of perfect tensors, for which the entanglement entropies are given exactly by the minimal cut through the underlying square lattice with randomly broken bonds.

Finally, for p = d = 2 we established that the purity of the reduced density matrix of a finite interval exhibits multifractal scaling at the transition. This implies different critical scaling exponents for the averaged and typical purity (the latter being equal to the prefactor of the logarithm of subsystem size in the entanglement entropy), and a continuous spectrum of critical exponents associated with the scaling of (noninteger moment averages of) the purity. The same features are expected to be present for (powers of) other primes p, while the corresponding exponents will vary with p for the symmetry reasons discussed in this paper.

Our approach could be generalized to other subgroups of the unitary group: formally, the main algebraic object that is needed would be the commutant of the subgroup. Once the commutant is known, most of the formulas derived in this paper carry over—including the definition of the Weingarten functions, etc. It would be interesting to investigate whether other universality classes of entanglement transitions can be obtained in this way. As in the Haar case, an analytic understanding of the replica limit of the statistical mechanics models for finite d remains a clear challenge for future work. One particularly intriguing aspect of our results is that while the statistical mechanics approach is certainly very powerful and allowed us to uncover the intricate dependence of the symmetry group on the prime number p, some aspects of Clifford circuits that are "simpler" (such as the ability to simulate them classically, or the flat entanglement spectrum of the reduced density matrix) are not yet particularly transparent in the current formulation of our framework. Those special properties of the Clifford group might possibly lead to some simplifications in the statistical mechanics formulation, which we leave for future work.

ACKNOWLEDGMENTS

We thank Utkarsh Agrawal, Xiao Chen, Sarang Gopalakrishnan, Michael Gullans, David Huse, Chao-Ming Jian, Vedika Khemani, Jed Pixley, Andrew Potter, Yi-Zhuang You, Justin Wilson, Zhi-Cheng Yang, and Aidan Zabalo for useful discussions and collaborations on related projects. We acknowledge support from the Air Force Office of Scientific Research under Grant No. FA9550-21-1-0123 (R.V.), and the Alfred P. Sloan Foundation through a Sloan Research Fellowship (R.V.). This work was supported by the Heising-Simons Foundation (Y.L. and M.P.A.F.), and by the Simons Collaboration on Ultra-Quantum Matter, which is a grant from the Simons Foundation (651440, M.P.A.F.), and was supported in part by the National Science Foundation under Grant No. DMR-1309667 (A.W.W.L.). Use was made of computational facilities purchased with funds from the National Science Foundation (CNS-1725797) and administered by the Center for Scientific Computing (CSC). The CSC is supported by the California NanoSystems Institute and the Materials Research Science and Engineering Center (MRSEC; NSF DMR-1720256) at UC Santa Barbara.

APPENDIX A: DESCRIPTION AND BASIC PROPERTIES OF THE COMMUTANT $\Sigma_{\rho}(p)$ FOR THE CLIFFORD CASE

The following statements (from Ref. [65]) lead to the full characterization of the commutant $\Sigma_Q(p)$, which is presented in statement (iv) below, together with its properties relevant for the construction of the Clifford stat mech model, summarized in (v)–(vii):

(i) (Stochastic) orthogonal group. For every $O \in \mathcal{O}_Q(p)$ the subspace

$$T_{O} := \{ (O\vec{\mu}, \vec{\mu}) : \vec{\mu} \in (\mathbf{Z}_{p})^{Q} \}$$
(A1)

is an element of $\Sigma_Q(p)$, and any operator $\hat{r}(T)$ as introduced in (23) is invertible if and only if *T* is of the form $T = T_O$ for some $O \in \mathcal{O}_Q(p)$. Clearly, it follows from (23) that the operator $\hat{r}(T_O)$ maps basis element $|\vec{\mu}\rangle$ into basis element $|O\vec{\mu}\rangle$. We denote this operator simply by $\hat{r}(O)$, and we write simply $\hat{r}(T_O) = \hat{r}(O)$.

(ii) Left and right multiplication by (stochastic) orthogonal group. For any subspace $T \in \Sigma_Q(p)$ as introduced in (23), and for any stochastic orthogonal matrix $O \in \mathcal{O}_Q(p)$, the objects

$$OT := \{ (O\vec{\nu}, \vec{\mu}) : (\nu, \mu) \in T \}$$
(A2)

and

$$TO := \{ (\vec{\nu}, O^t \vec{\mu}) : (\nu, \mu) \in T \}$$

are themselves again elements of $\Sigma_Q(p)$, i.e., $OT, TO \in \Sigma_Q(p)$.

(iii) Compatibility with $\hat{r}(T)$. The action with stochastic orthogonal matrices $O, O' \in \mathcal{O}_Q(p)$ on subspaces $T \in \Sigma_Q(p)$, defined in (ii) above, translates naturally into the action on the corresponding operator $\hat{r}(T)$:

$$\hat{r}(O)\hat{r}(T)\hat{r}(O') = \hat{r}(OT)\hat{r}(O') = \hat{r}(O)\hat{r}(TO') = \hat{r}(OTO').$$
(A3)

(iv) Double coset decomposition of commutant. Statement (iii) above shows that the commutant $\Sigma_Q(p)$ is a disjoint union of *double cosets* with respect to the left and right actions of the stochastic orthogonal group, i.e.,

$$\Sigma_{\mathcal{Q}}(p) = \mathcal{O}_{\mathcal{Q}}(p) \,\bar{T}_1 \,\mathcal{O}_{\mathcal{Q}}(p) \cup \dots \cup \mathcal{O}_{\mathcal{Q}}(p) \,\bar{T}_k \,\mathcal{O}_{\mathcal{Q}}(p),$$
(A4)

where $\bar{T}_1, \ldots, \bar{T}_k$ denote representatives of different double cosets. It can be shown [65] that $k \leq Q$. Here $\bar{T}_1 = \Delta$ is the "diagonal subspace"

$$\Delta := \{ (\vec{\mu}, \vec{\mu}) : \vec{\mu} \in (\mathbf{Z}_p)^Q \}, \tag{A5}$$

so that
$$\mathcal{O}_Q(p)\overline{T}_1\mathcal{O}_Q(p) = \mathcal{O}_Q(p)$$
.

(v) Choice of representatives for double cosets. The operators $\hat{r}(T)$ associated with elements T of the commutant that are not in the first double coset $\mathcal{O}_Q(p)\bar{T}_1\mathcal{O}_Q(p) = \mathcal{O}_Q(p)$ are not invertible. Representatives \bar{T}_a with $a = 2, \ldots, k$ can be chosen²¹ such that $\hat{r}(\bar{T}_a)$ are projectors. In particular, one can choose double coset representatives \bar{T}_a , $a = 2, 3, \ldots, k$ (and the notation \bar{T}_a denotes from now on that such a choice has been made) so that each is associated²² with a projector $P_{CSS(N_a)}$ possessing a so-called "defect subspace" $N_a \subseteq (\mathbf{Z}_p)^Q$ of dimension dim (N_a) , and

$$\hat{r}(\bar{T}_a) = p^{\dim(N_a)} P_{CSS(N_a)} \quad (a = 2, 3, \dots, k),$$
 (A6)

with trace

$$\operatorname{tr}[\hat{r}(\bar{T}_a)] = p^{Q-\dim(N_a)}.$$
(A7)

It is convenient to also introduce the notation $\bar{T}_1 = \Delta$ and $N_1 = 0$.

(vi) *Composition property.* Consider two arbitrary elements $T_1, T_2 \in \Sigma_Q(p)$ of the commutant. Each can then be reduced, upon right and left multiplication with elements in $\mathcal{O}_Q(p)$, to one of the elements²³ \overline{T}_a , where $a = 1, 2, \ldots, k$, and thus can be assigned a (possibly trivial) defect space N_a . Then the following composition property holds

$$\hat{r}(T_1)\hat{r}(T_2) = p^{\dim(N_1 \cap N_2)} \hat{r}(T_1 T_2).$$
(A8)

(vii) *Trace.* The commutant $\Sigma_Q(p)$ is equal to the disjoint union of sets $\Sigma_Q^{\ell}(p)$, where $\ell = 0, 1, ..., (Q-1)$, whose elements are those elements $T \in \Sigma_Q(p)$ with the property that

²¹Upon right and left multiplication with suitable stochastic orthogonal matrices, and making use of Theorem 4.24 of Ref. [65].

²²The subscript *CSS* stands for the so-called Calderbank-Shor-Steane codes [76–78] known from quantum information theory.

²³or to elements that are conjugate to those upon conjugation in $\mathcal{O}_Q(p)$, which then have again the same right and left defect spaces, which are images under $\mathcal{O}_Q(p)$ of the original ones.

 $\dim(T \cap \Delta) = Q - \ell$. Then, by the definition (23) of $\hat{r}(T)$ and the definition of the trace, one has²⁴

$$\operatorname{tr}[\hat{r}(T)] = p^{Q-\ell}, \text{ for all elements } T \in \Sigma_Q^{\ell}(p).$$
 (A9)

(viii) *Metric on an edge*. For any two elements of the commutant $\Sigma_Q(p)$, in analogy with (20) in the Haar case, the Boltzmann weight on an edge of the Clifford RTN is built from

$$\frac{1}{p^{\varrho}} \operatorname{tr}([\hat{r}(T_1)]^{\dagger} \hat{r}(T_2)) = e^{-\ln(p)|T_1, T_2|}, \quad (A10)$$

where is $|T_1, T_2|$ a metric defined by collecting (A6)–(A9). It satisfies the usual conditions $|T_1, T_2| = |T_2, T_1| \ge 0$, and $|T_1, T_2| = 0$ if and only if $T_1 = T_2$. We refer the reader to Appendix E for additional details.

As a check, let us briefly compute the metric in the case where $T_1 = T_2 = T$, and thereby confirm that the above rules indeed yield the expected result |T, T| = 0. There are two cases to consider. In the first case, $T \in \mathcal{O}_{\mathcal{O}}(p)$ is itself a group element, i.e., an element of the first double coset with "trivial" representative $T_1 = \Delta$. In this case the defect spaces are trivial (of dimension zero), and the trace in (A10) is that of the identity operator acting on the (replicated) edge Hilbert space $(\mathbf{C}^p)^Q$, yielding p^Q . This is canceled by the prefactor in (A10) with the result that |T, T| = 0 in this case, as expected. In the second case, T is an element of one of the remaining (k-1)double cosets and is thus, upon right and left multiplication with possibly different group elements²⁵ in $\mathcal{O}_{Q}(p)$, equal to one of the "projector" representatives \overline{T}_a with $a = 2, \dots, Q$. In this case $\bar{T}_a = P_{CSS(N_a)}$ is a projector satisfying $(\bar{T}_a)^2 = \bar{T}_a$, and (A6)-(A8) yield for (A10) the expression

$$\frac{1}{p^Q}p^{\dim(N_a)}p^{Q-\dim(N_a)}=1,$$

in agreement with the expected result |T, T| = 0.

(ix) Symmetry of the metric. It follows from its definition in (A10) and from the compatibility property (A3) that the metric $|T_1, T_2|$ is invariant under the symmetry group $\mathcal{O}_Q(p) \times \mathcal{O}_Q(p)$, i.e., under right and left multiplication with arbitrary elements of the stochastic orthogonal group,

$$|T_1, T_2| = |T_1', T_2'|$$
, where $T_1, T_2 \in \Sigma_Q(p)$, (A11)

and

$$T_1' = O_L T_1 O_R^{-1}, \quad T_2' = O_L T_2 O_R^{-1}, \text{ with } O_L, \quad O_R \in \mathcal{O}_Q(p).$$
(A12)

Just like in the Haar case, the metric is also invariant under a \mathbb{Z}_2 symmetry that corresponds to exchanging ket and bra. To see this, for each element of the commutant $T \in \Sigma_Q(p)$, we define T^{\dagger} as

$$\hat{r}(T) = \hat{r}(O_L)\hat{r}(\bar{T})\hat{r}(O_R) = \hat{r}(O_L\bar{T}O_R),$$
$$\hat{r}^{\dagger}(T) = \hat{r}(O_R^t)\hat{r}(\bar{T})\hat{r}(O_L^t) = \hat{r}(O_R^t\bar{T}O_L^t) := \hat{r}(T^{\dagger}),$$
where $\hat{r}^{\dagger}(\bar{T}) = \hat{r}(\bar{T}), \quad \bar{T}^{\dagger} = \bar{T}.$

Thus:
$$T = O_L \overline{T} O_R$$
, $T^{\dagger} := O_R^t \overline{T} O_L^t$

²⁴Recall (A5).

where we have used (A3), (A4), and (A6). With this notation in hand, we establish the following simple property of the metric:

$$|T_a, T_b| = |T_a^{\dagger}, T_b^{\dagger}|, \qquad (A13)$$

which follows from Eq. (A10) and

$$\operatorname{tr}\left(\left[\hat{r}(T_{a})\right]^{\dagger}\hat{r}(T_{b})\right) = \sum_{\left(\vec{v}_{a},\vec{\mu}_{a}\right)\in T_{a}}\sum_{\left(\vec{v}_{b},\vec{\mu}_{b}\right)\in T_{b}}\delta_{\vec{v}_{a},\vec{v}_{b}}\delta_{\vec{\mu}_{a},\vec{\mu}_{b}}$$
$$= \operatorname{tr}\left(\hat{r}(T_{a})\left[\hat{r}(T_{b})\right]^{\dagger}\right), \qquad (A14)$$

using Eq. (23). We note that Eq. (A14), in conjunction with (A10), also provides an explicit elementary definition of the metric.

It is straightforward to show that the Weingarten functions defined in Appendix D below also satisfy²⁶ $Wg_D(T_a^{\dagger}, T_b^{\dagger}) = Wg_D(T_a, T_b)$, which also implies that they are real.²⁷ Thus both principal ingredients of the Clifford statistical mechanics model for monitored circuits and RTNs,²⁸ namely, the Weingarten function $Wg_D(T_i, T_j)$ as well as the metric $|T_i, T_j|$, are invariant under the \mathbb{Z}_2 symmetry which sends $T_i \to T_i^{\dagger}$ at all sites.

APPENDIX B: BASIC PROPERTIES OF THE CLIFFORD GROUP, STABILIZER STATES, ETC.

(1) Single qudit (d = p = prime), Hilbert space $\mathcal{H}_1 = \mathbf{C}^p$, computational basis $|q\rangle$ with $q \in \{0, \dots, (p-1)\} = \mathbf{Z}_p$. *Pauli operators*

$$\begin{split} \hat{X}|q\rangle &= |q+1\rangle, \quad \hat{Z}|q\rangle = \omega^{q}|q\rangle, \ (\omega = e^{2\pi i q/d}), \\ \hat{Y} &:= \tau \hat{X}^{\dagger} \hat{Z}^{\dagger} \text{ (where } \tau = \text{some suitable phase),} \end{split}$$

then:
$$\hat{X}\hat{Y}\hat{Z} = \tau \mathbf{I}.$$
 (B1)

(2) *Hilbert space* for *n* qudits $(d = p = \text{prime}) \mathcal{H}_n = (\mathbf{C}^p)^n$, computational basis $|\mathbf{q}\rangle = (q_1, \ldots, q_n) \in (\mathbf{Z}_p)^n$.

(3) *Pauli group* \mathcal{P}_n is the group generated by the tensor product of all Pauli operators on the *n* qudits, or, equivalently, owing to (B1), by the group generated by the tensor product of the operators $\tau \hat{\mathbf{I}}, \hat{X}, \hat{Z}$.

(4) *Clifford group* Cliff(n, p) of the Hilbert space $\mathcal{H}_n = (\mathbb{C}^p)^n$ of *n* qudits is the normalizer of the Pauli group in the unitary group acting on $\mathcal{H}_n = (\mathbb{C}^p)^n$, i.e., the group of all unitary operators \hat{V} acting on $\mathcal{H}_n = (\mathbb{C}^p)^n$ satisfying $\hat{V}\mathcal{P}_n\hat{V}^{\dagger} \subset \mathcal{P}_n$ up to phases.

(5) For a subgroup of the Pauli group, $S \subset P_n$ [one that does not contain any (nontrivial) multiple of the identity operator], the operator

$$\hat{\boldsymbol{P}}_{\mathcal{S}} := \frac{1}{|\mathcal{S}|} \sum_{\hat{P} \in \mathcal{S}} \hat{P},$$

²⁵These group elements disappear when forming the trace in (A10).

²⁶Which follows from (D5) and the paragraph below this equation, upon making use of (A13).

²⁷Which follows from (D3) due to the manifest Hermiticity of \hat{Y} defined in (D1) (analogous to the Haar case as discussed in Appendix C).

 $^{^{28}}$ RTNs in the forced measurement formulation discussed in Sec. III E also require the use the Weingarten function.

is an orthogonal projector onto a subspace²⁹ $V_S \subset \mathcal{H}_n$ of dimension $p^n/|S|$. When $|S| = p^n$, the projection is onto a single state $|S\rangle$ called a *stabilizer state* (i.e., $V_S = \mathbf{C}|S\rangle$), which is thus the unique (up to scalars) eigenvector of all Pauli operators \hat{P} in S,

$$\hat{P}|\mathcal{S}\rangle = |\mathcal{S}\rangle$$
 (for all $\hat{P} \in \mathcal{S}$).

The (finite) set of all (pure) stabilizer states in $\mathcal{H}_n = (\mathbf{C}^p)^n$ is denoted Stab(n, p). The number of stabilizer states [79] in Stab(n, p) is $= p^n \prod_{i=1}^n (p^i + 1)$.

(6) For every stabilizer state $|S\rangle$ in $\mathcal{H}_n = (\mathbb{C}^p)^n$ there exists some Clifford unitary $\hat{V} \in \text{Cliff}(n, p)$ such that

$$|\mathcal{S}\rangle = \hat{V}(|0\rangle^{\otimes n})$$
 (where $|0\rangle = |q = 0\rangle$, with $q = 0 \in \mathbf{F}_p$).

It follows from this statement that the set of stabilizer states $\operatorname{Stab}(n, p)$ of the qudit Hilbert space $\mathcal{H}_n = (\mathbb{C}^p)^n$ is a *single orbit* of the action under the Clifford group $\operatorname{Cliff}(n, p)$.

(7) There is a simple way to see that the state $|0\rangle^{\otimes n} \in \mathcal{H}_n = (\mathbb{C}^p)^n$, is a stabilizer state, i.e., an element of the set Stab(n, p) of stabilizer states.

The argument uses the notation of Weyl operators:

$$\hat{W}_{\mathbf{x}} = \hat{W}_{\mathbf{p},\mathbf{q}} = \tau^{-\mathbf{p}\cdot\mathbf{q}}(\hat{Z}^{p_1}\hat{X}^{q_1}) \otimes \cdots \otimes (\hat{Z}^{p_n}\hat{X}^{q_n}),$$

where $\mathbf{x} = (\mathbf{p}, \mathbf{q}) \in \mathcal{V}_n := (\mathbf{Z}_p)^{2n}$ ("phase space"). Each Weyl operator is an element of the Paul group \mathcal{P}_n ; moreover, each element of the Pauli group is, up to a phase, equal to a Weyl operator. It is obvious [using (B1)] that the state $|0\rangle^{\otimes n} \in \mathcal{H}_n$ is the simultaneous eigenvector of the set of all Weyl operators $\hat{W}_{\mathbf{x}}$ with $\mathbf{x} = (\mathbf{p}, \mathbf{q})$ where $\mathbf{q} = \mathbf{0}$, i.e.,

$$\hat{W}_{\mathbf{n},\mathbf{0}}|0\rangle^{\otimes n} = |0\rangle^{\otimes n}.\tag{B2}$$

This particular set of these Weyl operators thus represents the subgroup $S \subset \mathcal{P}_n$ of the Pauli group, which "stabilizes" the state $|0\rangle^{\otimes n}$. Note that the order of this subgroup of the Pauli group is $|S| = p^n$, as required.

(8) Furthermore, it is now possible to show that $\hat{V}|0\rangle^{\otimes n}$, where $\hat{V} \in \text{Cliff}(n, p)$ is an arbitrary Clifford unitary operator in \mathcal{H}_n , is also a stabilizer state, i.e., an element of the set Stab(n, p) of stabilizer states. To see this, rewrite (B2) as

$$(\hat{V}\hat{W}_{\mathbf{p},\mathbf{0}}\hat{V}^{\dagger})(\hat{V}|0\rangle^{\otimes n}) = (\hat{V}|0\rangle^{\otimes n}),$$

which says that the group of p^n operators $(\hat{V}\hat{W}_{\mathbf{p},\mathbf{0}}\hat{V}^{\dagger})$ stabilizes the state $(\hat{V}|0\rangle^{\otimes n})$. Owing to Eq. (2.8) of Ref. [65] we know that

$$\hat{V}\hat{W}_{\mathbf{p},\mathbf{0}}\hat{V}^{\dagger} = \omega^{f_{\hat{V}}(\mathbf{p},\mathbf{0})} \hat{W}_{\Gamma_{\hat{V}}(\mathbf{p},\mathbf{0})}$$

where $\Gamma_{\hat{V}}(\mathbf{p}, \mathbf{0}) \in Sp(2n, p)$ and $f_{\hat{V}}(\mathbf{x})$ is a suitable function on phase space \mathcal{V}_n [defined under item (7) above—Sp(2n, p)is the symplectic group with entries in the field \mathbf{F}_p .] Then since, as also mentioned under item (7) above, every Weyl operator is an element of the Pauli group, the group $\{\hat{V}\hat{W}_{\mathbf{p},\mathbf{0}}\hat{V}^{\dagger}:$ $\mathbf{p} \in (\mathbf{Z}_p)^n\}$ forms a subgroup of the Pauli group \mathcal{P}_n and can be identified with the stabilizer subgroup of the stabilizer state $\hat{V}|0\rangle^{\otimes n}$. (9) Projective Clifford group, Pauli group, and orders of groups. Since we are interested in acting with elements of the Clifford group $\hat{V} \in \text{Cliff}(n, p)$ on a density matrix of a pure state $|\psi\rangle$, i.e., $\hat{V}|\psi\rangle \langle \psi|\hat{V}^{\dagger}$, we can work with elements of the Clifford group modulo phases. We define the projective Clifford group as the Clifford group modulo phases, i.e., $\overline{\text{Cliff}}(n, p) := \text{Cliff}(n, p)/U(1)$. Similarly, we call $\overline{\mathcal{P}}_n$ the Pauli group modulo phases. It can then be shown [65,79] that

$$\overline{\operatorname{Cliff}}(n, p)/\overline{\mathcal{P}}_n = \operatorname{Sp}_{2n}(p),$$

where, again, the latter is the symplectic group with entries in the field \mathbf{F}_p . Since the dimension of this symplectic group is known [79] to be $|\text{Sp}_{2n}(p)| = p^{n^2} \prod_{i=1}^n (p^{2i} - 1)$, we know that the order of the projective Clifford group is

$$|\overline{\text{Cliff}}(n,p)| = p^{2n} p^{n^2} \prod_{i=1}^n (p^{2i} - 1),$$
 (B3)

where we used the fact [see item (7) above] that the projective Pauli group is simply the set of all Weyl operators (of which there are as many as elements in phase space V_n , defined above).

(10) Average over the Clifford group. It turns out [65] that, for an *arbitrary* (meaning: not a stabilizer state) state $|\psi\rangle \in \mathcal{H}_n$, the average

$$\sum_{\hat{V} \in \text{Cliff}(n,p)} [\hat{V}|\psi\rangle \langle \psi|\hat{V}^{\dagger}]^{\otimes Q}, \tag{B4}$$

will, while still (obviously) expressible as a linear combination of the elements $\hat{R}(T)$ of the commutant, in general not simply be an equal-weight superposition of the latter, but may depend³⁰ on details of the state $|\psi\rangle$ itself. However, if we take $|\psi\rangle$ to be any *stabilizer state* $|\mathcal{S}_0\rangle \in \text{Stab}(n, p) \in \mathcal{H}_n$, the sum can be expressed as an equal-weight sum over the commutant, namely,

$$\frac{1}{\mathcal{N}_{\text{norm}}} \sum_{\hat{V} \in \text{Cliff}(n,p)} [\hat{V} | \mathcal{S}_0 \rangle \langle \mathcal{S}_0 | \hat{V}^{\dagger}]^{\otimes Q}$$
$$= \sum_{|\mathcal{S}) \in \text{Stab}(n,p)} [|\mathcal{S}\rangle \langle \mathcal{S}|]^{\otimes Q} = \frac{1}{Z_{n,p,Q}} \sum_{T \in \Sigma_Q(p)} \hat{R}(T), \text{ (B5)}$$

where \mathcal{N}_{norm} and

$$Z_{n,p,Q} = p^n \prod_{k=1}^{(Q-2)} (p^k + p^n).$$

are normalization factors. The first equality follows because the set Stab(n, p) of stabilizer states is [see item (6) above] a single orbit under the action of the Clifford group on stabilizer states.³¹

 $^{^{29}}V_S$ is called a "stabilizer code."

³⁰Certain linear combinations of expressions (B4) with different nonstabilizer states $|\psi\rangle$ can even be shown to represent Haar averages [65].

³¹If the first sum is over the projective Clifford group $\overline{\text{Clifft}}(n, p)$, which gives the same result up to possibly an overall multiplicative constant, it follows from (B3) that the normalization factor would in this case read $\overline{\mathcal{N}}_{\text{norm}} = p^n p^{n^2} \prod_{i=1}^n (p^i - 1)$ because [79] every

APPENDIX C: WEINGARTEN FUNCTIONS FOR HAAR

Let us consider the operator³² (repeated indices summed)

x + + + × 0

$$\begin{split} X &:= \mathbb{E}_{\hat{U} \in U(D)} [U(\cdots)U^{\dagger}]^{\otimes \mathcal{Q}} \\ &= \int_{\text{Haar}} d\mu(\hat{U}) [\hat{U}(\cdots)\hat{U}^{\dagger}]^{\otimes \mathcal{Q}} \\ &= \int_{\text{Haar}} d\mu(U) U_{\tilde{i}_{1}, j_{1}} \cdots U_{\tilde{i}_{\mathcal{Q}}, j_{\mathcal{Q}}} U_{\tilde{j}_{1}, i_{1}}^{\dagger} \cdots U_{\tilde{j}_{\mathcal{Q}}, i_{\mathcal{Q}}}^{\dagger} \\ &\times (|\tilde{i}_{1}, \dots, \tilde{i}_{\mathcal{Q}}\rangle \langle i_{1}, \dots, i_{\mathcal{Q}}|) \otimes (|\tilde{j}_{1}, \dots, \tilde{j}_{\mathcal{Q}}\rangle \langle j_{1}, \dots, j_{\mathcal{Q}}|), \end{split}$$

where $i_1, \ldots, \bar{i}_Q, j_1, \ldots, \bar{j}_Q \in \{1, \ldots, D\}$. We can view the second tensor factor in the fourth line, $(|\bar{j}_1, \ldots, \bar{j}_Q\rangle\langle j_1, \ldots, j_Q|)$, as defining an orthonormal basis of the *Q*-fold tensor product of an "in" vector space of operators \mathcal{H}_{in}^{op} , i.e., of $(\mathcal{H}_{in}^{op})^{\otimes Q}$. In the same way, we can view the first tensor factor in the fourth line, $(|\bar{i}_1, \ldots, \bar{i}_Q\rangle\langle i_1, \ldots, i_Q|)$, as defining an orthonormal basis of the *Q*-fold tensor product an "out" vector space of operators \mathcal{H}_{out}^{op} , i.e., of $(\mathcal{H}_{out}^{op})^{\otimes Q}$. In other words, we can view \hat{X} as an element of the tensor product of these two vector spaces of operators:

$$\hat{X} \in \left(\mathcal{H}_{\text{out}}^{\text{op}}\right)^{\otimes \mathcal{Q}} \otimes \left(\mathcal{H}_{\text{in}}^{\text{op}}\right)^{\otimes \mathcal{Q}}.$$
(C2)

Let us now consider a unitary operator \hat{W}_1 acting (by conjugation) solely on \mathcal{H}_{out} , while we act with the identity operator I on \mathcal{H}_{in} . For such an operator we obtain³³

$$\left(\hat{W}_{1}^{\otimes Q} \otimes \mathbf{I}^{\otimes Q}\right) \hat{X} \left(\hat{W}_{1}^{\dagger \otimes Q} \otimes \mathbf{I}^{\otimes Q}\right) = \hat{X}, \quad (C3)$$

which follows from the invariance of the Haar measure under left-multiplication of \hat{U} with \hat{W}_1 . Similarly, for a unitary operator \hat{W}_2 acting (by conjugation) solely on \mathcal{H}_{in} , while we act with the identity operator on \mathcal{H}_{out} , we obtain

$$\left(\mathbf{I}^{\otimes \mathcal{Q}} \otimes \hat{W}_{2}^{\dagger \otimes \mathcal{Q}}\right) \hat{X} \left(\mathbf{I}^{\otimes \mathcal{Q}} \otimes \hat{W}_{2}^{\otimes \mathcal{Q}}\right) = \hat{X}$$
(C4)

due to invariance of the Haar measure under rightmultiplication of \hat{U} by \hat{W}_2 . Owing to (C3) and (C4), the operator \hat{X} can be expanded as a tensor product of operators $\hat{R}(\sigma)$ which form a basis of the commutant acting on $(\mathcal{H}_{in}^{op})^{\otimes Q}$, and operators $\hat{R}(\tau)$ which form a basis of the commutant acting on $(\mathcal{H}_{out}^{op})^{\otimes Q}$, i.e.,

$$\hat{X} = \sum_{\sigma, \tau \in S_Q} Wg_D(\sigma, \tau) \hat{R}(\sigma) \otimes \hat{R}(\tau),$$
(C5)

³³More explicitly, the left-hand side is the Haar average of $U_{\tilde{i}_1,j_1}\cdots U_{\tilde{i}_Q,j_Q}U_{\tilde{j}_1,i_1}^{\dagger}\cdots U_{\tilde{j}_Q,i_Q}^{\dagger}(\hat{W}_1^{\otimes Q}|\tilde{i}_1,\ldots,\tilde{i}_Q\rangle\langle i_1,\ldots,i_Q|\hat{W}_1^{\dagger\otimes Q}) \otimes (|\tilde{j}_1,\ldots,\tilde{j}_Q\rangle\langle j_1,\ldots,j_Q|),$ equal to \hat{X} upon letting $U \to W_1U$.

where $Wg_D(\sigma, \tau)$ are expansion coefficients³⁴ called Weingarten functions. Clearly, reading (C1) in conjunction with (C5), (17), and (18), equation (C1) is equivalent to the statement (10) and constitutes a proof of the latter.

We close this discussion with a slightly more compact, basis-independent formulation of the same facts. In particular, if we apply \hat{X} to an operator in $(\mathcal{H}_{in}^{op})^{\otimes Q}$ that is a *Q*-fold tensor product of an operator \hat{O}_{in} which is an element in the Hilbert space \mathcal{H}_{in}^{op} of in operators, we obtain the following result which is an element of the *Q*-fold tensor product in the Hilbert space \mathcal{H}_{opt}^{op} of out operators:

$$\hat{X} : \left(\mathcal{H}_{in}^{op}\right)^{\otimes Q} \to \left(\mathcal{H}_{out}^{op}\right)^{\otimes Q}, \\
(\hat{O}_{in})^{\otimes Q} \to \hat{X}[(\hat{O}_{in})^{\otimes Q}],$$
(C6)

where

$$\hat{X}[(\hat{O}_{\rm in})^{\otimes Q}] = \int_{\rm Haar} d\mu(\hat{U})[\hat{U}\hat{O}_{\rm in}\hat{U}^{\dagger}]^{\otimes Q}.$$

The operator \hat{X} is (evidently) invariant under the action of independent unitary operators \hat{W}_1 and \hat{W}_2 , implemented by right and left multiplication of the (integrated) Haar unitary \hat{U}

$$\begin{split} \hat{X}[(\hat{O}_{\mathrm{in}})^{\otimes \mathcal{Q}}] &= \hat{X}\left[\hat{W}_{2}^{\otimes \mathcal{Q}}\hat{O}_{\mathrm{in}}^{\otimes \mathcal{Q}}\hat{W}_{2}^{\dagger \otimes \mathcal{Q}}\right] \\ &= \hat{W}_{\mathrm{l}}^{\otimes \mathcal{Q}}(\hat{X}[(\hat{O}_{\mathrm{in}})^{\otimes \mathcal{Q}}])\hat{W}_{\mathrm{l}}^{\dagger \otimes \mathcal{Q}} \end{split}$$

This is the symmetry giving rise to the expansion (C5) of \hat{X} in terms of a tensor product of elements of the commutant. The same logic will apply in the Clifford case, the only difference being that the commutant will then be different.

We note for future use the fact that we can apply the operator \hat{X} to any element $\hat{R}(\tau)$ of the commutant, viewed as an element of the vector space of operators $(\mathcal{H}_{in}^{op})^{\otimes Q}$. Using the fact that $\hat{R}(\tau)$ commutes by definition with $\hat{U}^{\otimes Q}$, we see that $\hat{X}[\hat{R}(\tau)] = \hat{R}(\tau)$ where, by (C6), the right-hand side is an element of the vector space $(\mathcal{H}_{out}^{op})^{\otimes Q}$.

1. Key properties of the Weingarten functions $Wg_D(\sigma, \tau)$ for the Haar case

Let us perform the trace over the second tensor factor $(\mathcal{H}_{in}^{op})^{\otimes Q}$ of \hat{X} defined in (C1) and (C2); in the following we denote the trace over the first³⁵ and second³⁶ tensor factor by tr⁽¹⁾ and tr⁽²⁾. When denoting the identity operator in the Hilbert space underlying the first tensor factor, i.e., in \mathcal{H}_{out} , by **I** we obtain for any permutation $\tau \in S_Q$

$$\begin{aligned} \operatorname{tr}^{(2)} \hat{X}(\mathbf{I}^{\otimes \mathcal{Q}} \otimes \hat{R}(\tau)) &= \hat{X}[\hat{R}(\tau)] = \hat{R}(\tau) \\ &= \sum_{\sigma_o, \sigma_i \in S_{\mathcal{Q}}} \operatorname{Wg}_D(\sigma_o, \sigma_i) \hat{R}(\sigma_o) \operatorname{tr}[\hat{R}^{\dagger}(\sigma_i) \hat{R}(\tau)] \\ &= \sum_{\sigma_o, \sigma_i \in S_{\mathcal{Q}}} \operatorname{Wg}_D(\sigma_o, \sigma_i) \hat{R}(\sigma_o) D^{\mathcal{Q} - |\sigma_i, \tau|} \in \left(\mathcal{H}_{\operatorname{out}}^{\operatorname{op}}\right)^{\otimes \mathcal{Q}}, \quad (C7) \end{aligned}$$

³⁴Which are real because \hat{X} is Hermitian, and $Wg_D(\sigma, \tau) = Wg_D(\sigma^{-1}, \tau^{-1})$.

³⁵In other words, over $(\mathcal{H}_{out}^{op})^{\otimes Q}$. ³⁶In other words, over $(\mathcal{H}_{in}^{op})^{\otimes Q}$.

element of $\overline{\text{Cliff}}(n, p)$ is a product of an element of the Pauli group (modulo phases) and a (projective) representation of the symplectic group $Sp_{2n}(p)$. The result follows because a subgroup of order p^n of the Pauli group is the stabilizer group of the stabilizer state, and the length of the orbit is $|\text{Stab}(n, p)| = p^n \prod_{i=1}^n (p^i + 1)$ (as mentioned above).

³²The last two lines are obtained from the second line by inserting a complete set of *D* states in each of the *Q* tensor factors, e.g., $\hat{U} = |\vec{i}_1\rangle U_{\vec{i}_1,\vec{j}_1} \langle j_1|$, etc.—we use the notation $|\vec{i}_1, \ldots, \vec{i}_Q\rangle = |\vec{i}_1\rangle \otimes \ldots \otimes |\vec{i}_Q\rangle$, etc.

where in the first line we used the fact that $\hat{R}(\tau)$ is an element of the commutant [and the definition (C1) of \hat{X} , as well as of the metric, (22) and (21)], and in the second line we used (C5). Here $D = d^2$ is the dimension of the Hilbert space on which the two-site unitary gate acts. Since $\hat{R}(\sigma_o)$ form a basis of the commutant, (C7) implies that

$$\sum_{\sigma_i} Wg_D(\sigma_o, \sigma_i) D^{Q-|\sigma_i, \tau|} = \delta_{\sigma_o, \tau}.$$
 (C8)

This equation says that the Weingarten function is the leftinverse of $D^{Q-|\sigma_i,\tau|}$ under matrix multiplication when both are viewed as matrices.

The analogous argument applied to the equation $\operatorname{tr}^{(1)} \hat{X}(\hat{R}(\tau) \otimes \mathbf{I}^{\otimes Q}) = \hat{R}(\tau)$ implies a similar equation for the right inverse,

$$\sum_{\sigma_o} D^{\mathcal{Q}-|\tau,\sigma_o|} Wg_D(\sigma_o,\sigma_i) = \delta_{\tau,\sigma_i}.$$
 (C9)

A simplification occurs in the Haar case because the metric satisfies $|\sigma, \tau| = |\sigma^{-1}\tau|$, i.e., it does not depend separately on σ and τ . This has two implications: (i) First, and most importantly, the Weingarten function, being the inverse of $D^{Q-|\sigma,\tau|} = D^{Q-|\sigma^{-1}\tau|}$ is also a function of the combined variable, namely, $Wg_D(\sigma_o, \sigma_i) = Wg_D(\sigma_o^{-1}\sigma_i)$. (ii) Second, without loss of generality, we may replace $\tau \to 1$ by the identity permutation 1 in (C8) and (C9).

2. Consequences of (9) and (11)

Set $j_1 = \cdots = j_Q = \overline{j}_1 = \cdots = \overline{j}_Q = 1$ in those equations. If we define the projection operator onto this state in $(\mathcal{H}_{in})^{\otimes Q}$,

$$\hat{\mathbf{Q}}^0 := |1\rangle^{\otimes \mathcal{Q}} \langle 1|^{\otimes \mathcal{Q}},$$

we conclude that

$$\begin{aligned} \mathrm{tr}^{(2)}[\hat{X} (\mathbf{I}^{\otimes Q} \otimes \hat{\mathbf{Q}}^{0})] &= \int_{\mathrm{Haar}} d\mu(\hat{U})[\hat{U}|1\rangle \langle 1|\hat{U}^{\dagger}]^{\otimes Q} \\ &= \mathrm{const} \times \sum_{\tau \in S_{Q}} \hat{R}(\tau), \end{aligned}$$

where we used (9) and (11).

On the other hand, using the form (C5) for \hat{X} , the same expression is equal to

$$= \sum_{\sigma,\tau \in S_Q} \operatorname{Wg}_D(\sigma,\tau) \hat{R}(\sigma)(\operatorname{tr}[\hat{R}(\tau)\hat{\mathbf{Q}}^0]).$$

Owing to (12) the trace in the above equation is unity independent³⁷ of the element τ of the commutant S_Q ,

$$(\operatorname{tr}[\hat{R}(\tau) \, \hat{\mathbf{Q}}^0]) = 1 \quad \text{for all } \tau \in S_Q.$$

We therefore conclude that

$$\operatorname{const} \times \sum_{\sigma \in S_Q} \hat{R}(\sigma) = \sum_{\sigma \in S_Q} \left(\sum_{\tau \in S_Q} \operatorname{Wg}_D(\sigma, \tau) \right) \hat{R}(\sigma),$$

³⁷Because the state $|1\rangle^{\otimes Q}$ is invariant under permutation of the *Q* tensor factors.

which means that the expression in parentheses is a constant, i.e., independent of σ ,

$$\sum_{\tau \in S_Q} Wg_D(\sigma, \tau) = \text{const} \quad \text{for all } \sigma \in S_Q.$$
 (C10)

Of course, here in the Weingarten case for Haar under consideration here, $Wg_D(\sigma, \tau)$ is a function of only $\sigma \tau^{-1}$ and so this condition is obviously satisfied.

APPENDIX D: WEINGARTEN FUNCTIONS FOR CLIFFORD

Let us start with the analog of (C1) for Clifford:

$$\begin{split} \hat{Y} &:= \mathbb{E}_{\hat{V} \in \operatorname{Cliff}(n,p)} [\hat{V}(\cdots)\hat{V}^{\dagger}]^{\otimes Q} \\ &= \frac{1}{\mathcal{N}_{\operatorname{Cliff}}} \sum_{\hat{V} \in \operatorname{Cliff}(n,p)} [\hat{V}(\cdots)\hat{V}^{\dagger}]^{\otimes Q} \\ &= \frac{1}{\mathcal{N}_{\operatorname{Cliff}}} \sum_{\hat{V} \in \operatorname{Cliff}(n,p)} V_{\bar{i}_{1},j_{1}} \cdots V_{\bar{i}_{Q},j_{Q}} V_{\bar{j}_{1},i_{1}}^{\dagger} \cdots V_{\bar{j}_{Q},i_{Q}} \\ &\times (|\bar{i}_{1},\ldots,\bar{i}_{Q}\rangle\langle i_{1},\ldots,i_{Q}|) \otimes (|\bar{j}_{1},\ldots,\bar{j}_{Q}\rangle\langle j_{1},\ldots,j_{Q}|), \end{split}$$
(D1)

where

$$\hat{Y} \in \left(\mathcal{H}_{\text{out}}^{\text{op}}\right)^{\otimes Q} \otimes \left(\mathcal{H}_{\text{in}}^{\text{op}}\right)^{\otimes Q}.$$
 (D2)

Note that here $|i\rangle \in \mathcal{H}_n = (\mathbb{C}_p)^n$ denotes elements of an *or*thonormal basis of \mathcal{H}_n , having no relationship with stabilizer states in \mathcal{H}_n (but see Sec. D 2 below). Now, the analogs of (C3) and (C4) follow up on the replacement $\hat{X} \to \hat{Y}$, for any *Clifford* unitary operators \hat{W}_1 and \hat{W}_2 . Furthermore, the analog of (C5) for the expansion of \hat{Y} as a tensor product of operators in the commutant goes through in the same way as for Haar, leading to

$$\hat{Y} = \sum_{T_o, T_i \in \Sigma_Q(p)} Wg_D(T_o, T_i)\hat{R}(T_o) \otimes \hat{R}(T_i),$$
(D3)

where $Wg_D(T_o, T_i)$ are generalized Weingarten functions (see also Ref. [80] for explicit expressions in the case of Q = 4 replicas).

1. Key properties of the Weingarten functions $Wg_D(T_o, T_i)$ for the Clifford case

We proceed in complete analogy with the Haar case, discussed in Sec. C 1. For any element of the commutant $T \in \Sigma_Q(p)$, we have

$$\operatorname{tr}^{(2)} \hat{Y}(\mathbf{I}^{\otimes Q} \otimes \hat{R}(T)) = \hat{R}(T)$$

$$= \sum_{T_o, T_i \in \Sigma_Q(p)} \operatorname{Wg}_D(T_o, T_i) \hat{R}(T_o) \operatorname{tr}[\hat{R}^{\dagger}(T_i) \hat{R}(T)]$$

$$= \sum_{T_o, T_i \in \Sigma_Q(p)} \operatorname{Wg}_D(T_o, T_i) \hat{R}(T_o) D^{Q-|T_i, T|}, \quad (D4)$$

where in the first line we used the fact that T belongs to the commutant. We thus have

$$\sum_{T_i \in \Sigma_Q(p)} Wg_D(T_o, T_i) D^{Q-|T_i, T|} = \delta_{T_o, T}.$$
 (D5)

The last equation says that the Weingarten function is the leftinverse of $D^{Q-|T_i,T|}$ (both viewed as a matrix). Similarly, by acting in the first line of (D4) with $\hat{R}(T)$ on the first tensor factor and performing the trace over the first, we get a similar equation for the right inverse.

2. Connection with stabilizer states—consequences of (B5)

To this end, we pick an orthonormal basis in $(\mathbf{C}^p)^n$ so that one element of this basis is the state $|0\rangle^{\otimes n} \in \mathcal{H}_n = (\mathbf{C}^p)^n$ where 0 denotes the number zero in the finite number field \mathbf{F}_p ; this is the state that is described in items (6) and (7) of Appendix B. Having done so, we use this orthonormal basis in (D1) and set $j_k = j_k^0$, $\bar{j}_k = \bar{j}_k^0$ for $k = 1, \ldots, Q$, where $j_1^0 = \cdots = j_Q^0 = \bar{j}_1^0 = \cdots = \bar{j}_Q^0 = 0^{\otimes n}$. (In better notation, $|j_1^0\rangle = \cdots = |j_Q^0\rangle = |0\rangle^{\otimes n}$, etc.)

If we define the projection operator onto this state in $(\mathcal{H}_{in})^{\otimes \mathcal{Q}}$

$$\hat{\mathbf{P}}^0 := [|0\rangle^{\otimes n}]^{\otimes Q} [\langle 0|^{\otimes n}]^{\otimes Q},$$

we conclude that

$$\begin{aligned} & \operatorname{tr}^{(2)}[\hat{Y} \ (\mathbf{I}^{\otimes Q} \otimes \hat{\mathbf{P}}^{0})] \\ &= \frac{1}{\mathcal{N}_{\operatorname{Cliff}}} \sum_{\hat{V} \in \operatorname{Cliff}(n,p)} [\hat{V}(|0\rangle^{\otimes n})(\langle 0|^{\otimes n})\hat{V}^{\dagger}]^{\otimes Q} \\ &= \operatorname{const} \times \sum_{T \in \Sigma_{Q}(p)} \hat{R}(T), \end{aligned}$$

where we used (B5).

On the other hand, using the form (D3) for \hat{Y} , the same expression is equal to

$$= \sum_{T_o, T_i \in \Sigma_Q(p)} \operatorname{Wg}_D(T_o, T_i) \hat{R}(T_o)(\operatorname{tr}[\hat{R}(T_i) \, \hat{\mathbf{P}}^0]).$$

Owing to Eq. (4.10) of Ref. [65], the trace in the above equation is unity independent of the element T_i of the commutant $\Sigma_Q(p)$,

$$(\operatorname{tr}[\hat{R}(T_i)\hat{\mathbf{P}}^0]) = 1 \text{ for all } T_i \in \Sigma_Q(p).$$

We therefore conclude that

$$\operatorname{const} \times \sum_{T \in \Sigma_{\mathcal{Q}}(p)} \hat{R}(T)$$
$$= \sum_{T_o \in \Sigma_{\mathcal{Q}}(p)} \left(\sum_{T_i \in \Sigma_{\mathcal{Q}}(p)} \operatorname{Wg}_D(T_o, T_i) \right) \hat{R}(T_o),$$

which means that the expression in parentheses is a constant, independent of T_o ,

$$\sum_{T_i \in \Sigma_Q(p)} Wg_D(T_o, T_i) = \text{const} \quad \text{for all } T_o \in \Sigma_Q(p).$$
 (D6)

This is the analog of (C10) of the Haar case, where it is tied to the group structure of the commutant.

3. Generalization to other groups

Note that the generalized Weingarten formulas of the previous section do not rely on any specific property of the commutant of the Clifford group. As such, most of our results can be generalized to derive statistical mechanics models for RTNs and monitored quantum circuits involving averages over other subgroups of the unitary group, expressed solely in terms of the commutant of such subgroups. Once this algebraic object, the commutant, is known, the link and Weingarten weights derived for the Clifford group carry over without any change, with the spins T_i now belonging to this new commutant. Previously, Weingarten functions had been derived for unitary [67], orthogonal and symplectic groups [81,82]—See Ref. [83] for a recent review. The formulas in the present Appendix are completely general, and extend the notion of the Weingarten function to all situations for which the commutant is known.

APPENDIX E: METRIC ON THE COMMUTANT OF THE CLIFFORD GROUP

The Boltzmann weights of the Clifford stat mech model rely on the existence of a *metric* on the commutant: for any two elements T_a and T_b of the commutant $\Sigma_Q(p)$, there exists a *metric* which arises from the trace:

$$W(T_a, T_b) := \operatorname{tr}[\hat{R}^{\dagger}(T_a)\hat{R}(T_b)] = (p^N)^{Q - |T_a, T_b|}, \quad (E1)$$

for any $T_a, T_b \in \Sigma_Q(p)$, where $\hat{R}(T)$ is defined in (25). Our main results rely only on the standard properties satisfied by this metric: $|T_a, T_b| \ge 0$, $|T_a, T_b| = |T_b, T_a|$, and $|T_a, T_b| = 0$ if and only if $T_a = T_b$. In this Appendix, we provide additional details on the definition of this metric following closely Ref. [65].

Consider the double coset representation of the commutant defined in Eq. (A4). The simplest elements are the orthogonal group elements themselves, $T_1 = O_1$, $T_2 = O_2 \in \mathcal{O}_Q(p)$, which, as noted in the line below (A5), correspond to the first element of the disjoint union of double cosets in (A4): When applied to those, \hat{r} is a representation of the group $\mathcal{O}_Q(p)$ [see our Eq. (A1)], and thus we have Eq. (A10). When $O_1 = O_2$, we have $\langle \vec{\mu} | (O'_1 O_2) | \vec{\mu} \rangle = 1$ for all $\vec{\mu}$, thus $e^{-\ln(p) |O_1, O_2|} = 1$ and $|O_1, O_2| = 0$. When $O_1 \neq O_2$, we have $\langle \vec{\mu} | (O'_1 O_2) | \vec{\mu} \rangle < 1$ for at least some $\vec{\mu}$, thus $e^{-\ln(p) |O_1, O_2|} < 1$ and $|O_1, O_2| > 0$.

Now consider two general elements T_1 , T_2 of the commutant $\Sigma_Q(p)$ (which may or may not be elements of the same double coset). According to Proposition 4.17, as well as Eq. (4.18) and Lemma 4.18 of Ref. [65], we can transform those by right and left multiplication with group elements in $\mathcal{O}_Q(p)$ to elements T'_1 and T'_1 ,

$$T_j \to T'_j \equiv O_j T_j O'_j, \quad \text{with} \quad O_j, O'_j \in \mathcal{O}_Q(p) \ (j = 1, 2),$$

so that the right and left defect subspaces $T'_{j,LD}$ and $T'_{j,RD}$ of each T'_{j} become equal, $N'_{j} \equiv T'_{j,LD} = T'_{j,RD}$, i.e.,

$$O_{j} T_{j,LD} = O_{j}^{\prime t} T_{j,RD} \ (j = 1, 2).$$
 (E2)

Since the last equality in Eq. (E2) above can be achieved by choosing³⁸ for example $O_1 = O_2 = \mathbf{1}$, let us make that choice so that

$$T'_1 = T_1 \mathcal{O}'_1, \quad T'_2 = T_2 \mathcal{O}'_2.$$
 (E3)

³⁸Letting $O_j \rightarrow \mathbf{1}$, we have $O'_j^t \rightarrow O_j^t O'_j^t = (O'_j O_j)^t$.

Then, using Eq. (A3) we get for the trace in (A10)

$$tr([\hat{r}(T_1)]^{\dagger}\hat{r}(T_2)) = tr(\hat{r}(O'_1)[\hat{r}(T'_1)]^{\dagger}\hat{r}(T'_2)\hat{r}(O'_2))$$

= tr($\hat{r}(O'_2 O'_1)[\hat{r}(T'_1)]^{\dagger}\hat{r}(T'_2)$). (E4)

Now, owing to (E2) above and Theorem 4.24 of Ref. [65], the operators $\hat{r}(T'_j)$ are both proportional to Calderbank-Shor-Steane (CSS) projectors,

$$\hat{r}(T'_j) = p^{\dim N'_j} P_{\mathrm{CSS}(N'_j)}.$$

Note in particular their Hermiticity, especially $[\hat{r}(T'_1)]^{\dagger} = \hat{r}(T'_1)$. Next, Eq. (4.24) of Ref. [65] proves the result

$$[\hat{r}(T'_1)]^{\dagger}\hat{r}(T'_2) = \hat{r}(T'_1)\hat{r}(T'_2) = p^{\dim(N'_1 \cap N'_2)}\hat{r}(T'_1T'_2),$$
(E5)

which (at the same time) defines the meaning of the product of T'_1 and T'_2 on the right-hand side (the "semigroup property").

Using these results, the trace in Eq. (E4) above then leads to the following expression for the metric $e^{-\ln(p)|T_1,T_2|}$:

$$e^{-\ln(p)[Q-\dim(N'_1\cap N'_2)]} \operatorname{tr}[\hat{r}(O'_2{}^tO'_1)\hat{r}(T'_1T'_2)].$$
(E6)

Note that in the special case where $T_1 = T'_1$ and $T_2 = T'_2$, i.e., when both are already proportional to (CSS) projectors, thus when both already have equal right and left defect subspaces, we also have $O'_j = \mathbf{1}$ for j = 1, 2 (see, e.g., Eq. (E3)]. In this case, using Lemma 4.25 of Ref. [65], we find³⁹

$$|T_1, T_2| = [Q - \dim(N_1 \cap N_2) - \dim(N_1^{\perp} \cap N_2^{\perp})],$$

and we have $|T_1, T_2| \ge 0$ since $(N_1 \cap N_2)^{\perp} \supseteq N_1^{\perp} + N_2^{\perp} \supseteq N_1^{\perp} \cap N_2^{\perp}$.

In the general case, Eq. (E6) above, where $\hat{r}(O'_2O'_1)$ is not the identity, a closed form expression is presently not known, even though an explicit elementary expression for the metric is given by (A14) and (A10), as already mentioned below (A14). Finally, we note that since the permutation group S_Q is a subgroup⁴⁰ of $\mathcal{O}_Q(p)$, it follows from its definition that the metric between two permutations is the same as that familiar from the Haar circuits and expressible in terms of the cyclecounting function, a property used in the main text.

APPENDIX F: SOME DETAILS ON MULTIFRACTAL SCALING

In this Appendix we present more details on the multifractal scaling of the purity, discussed in Sec. IV C. This is a special case of the general discussion of the scaling of a correlation function (at disorder dominated critical points) in disordered systems, as discussed in Refs. [49,72,73].

We start from the scaling of all moments of the purity at the transition, Eq. (60),

$$\overline{\left[G_{(x_1,x_2)}\right]^k} \sim \frac{B_k}{R_{12}^{2X_k}},$$
(F1)

where the purity in a fixed realization of circuit disorder is a random variable, viewed as a two-point correlation function $G(x_1, x_2) = \text{tr}(\hat{\rho}_{A,\mathbf{m}})^2$ associated with the boundary-condition changing (bcc) operator determining entanglement properties [25,56]. Here R_{12} is the chord distance:

$$R_{12} := \frac{L}{\pi} \sin\left(\frac{\pi}{L}|x_1 - x_2|\right) = \frac{L}{\pi} \sin\left(\frac{\pi}{L}|A|\right).$$

Since upon recalling (59), $\ln G(x_1, x_2) = -S_A^{(2)}$, the cumulant expansion of the left-hand side of (F1) reads

$$\overline{\left[G_{(x_1,x_2)}\right]^k} = \exp\left\{-k \,\overline{S_A^{(2)}} + \frac{k^2}{2!} \,\overline{\left(S_A^{(2)} - \overline{S_A^{(2)}}\right)^2} - \frac{k^3}{3!} \,\kappa_3[S_A^{(2)}] + \cdots\right\},\tag{F2}$$

where $\kappa_3[S_A^{(2)}]$ denotes the third cumulant of the random variable $S_A^{(2)}$. We compare this with the right-hand side of (F1), using the Taylor expansion (64) of the exponents X_k in powers of k,

$$X_k = k x^{(1)} + \frac{k^2}{2!} x^{(2)} + \frac{k^3}{3!} x^{(3)} + \cdots,$$
 (F3)

yielding

$$B_{k} \exp \{-2X_{k} \ln R_{12}\} = B_{k} \exp \left\{-\left[k2x^{(1)} + \frac{k^{2}}{2!}2x^{(2)} + \frac{k^{3}}{3!}2x^{(3)} + \cdots\right] \ln R_{12}\right\}.$$
(F4)

Comparison of (F2) with (F4) shows that all cumulants grow⁴¹ proportional to $\ln R_{12}$, the constants of proportionality being universal and equal to $(-1)^{k-1}$ times twice the expansion coefficients $x^{(k)}$ appearing the Taylor expansion (F3),

$$\overline{S_A^{(2)}} \sim 2x^{(1)} \ln R_{12},$$
 (F5)

$$\overline{\left(S_A^{(2)} - \overline{S_A^{(2)}}\right)^2} \sim -2x^{(2)} \ln R_{12},$$
 (F6)

$$\kappa_3[S_A^{(2)}] \sim 2x^{(3)} \ln R_{12},$$

:. (F7)

These are Eqs. (61)–(63) of the main text.

³⁹In particular, using the unnumbered equation below Eq. (4.26) of Ref. [65].

⁴⁰Recall, e.g., (24).

⁴¹The analogous expansion of $B_k = \exp\{k \ b_1 + (k^2/2!) \ b_2 + \cdots\}$ shows that a nonuniversal constant offset $(-1)^k \ b_k$ appears in the *k*th cumulant $\kappa_k[S_A^{(2)}]$ of the entanglement entropy $S_A^{(2)}$, but these offsets will of course just represent subleading contributions in the regime of interest of large values of R_{12} .

- L. Amico, R. Fazio, A. Osterloh, and V. Vedral, Entanglement in many-body systems, Rev. Mod. Phys. 80, 517 (2008).
- [2] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, Rev. Mod. Phys. 81, 865 (2009).
- [3] J. Eisert, M. Cramer, and M. B. Plenio, Colloquium: Area laws for the entanglement entropy, Rev. Mod. Phys. 82, 277 (2010).
- [4] P. Calabrese and J. Cardy, Entanglement entropy and conformal field theory, J. Phys. A: Math. Theor. 42, 504005 (2009).
- [5] N. Laflorencie, Quantum entanglement in condensed matter systems, Phys. Rep. 646, 1 (2016).
- [6] J. Preskill, Quantum Computing in the NISQ era and beyond, Quantum **2**, 79 (2018).
- [7] A. Nahum, J. Ruhman, S. Vijay, and J. Haah, Quantum entanglement growth under random unitary dynamics, Phys. Rev. X 7, 031016 (2017).
- [8] A. Nahum, S. Vijay, and J. Haah, Operator spreading in random unitary circuits, Phys. Rev. X 8, 021014 (2018).
- [9] C. W. von Keyserlingk, T. Rakovszky, F. Pollmann, and S. L. Sondhi, Operator hydrodynamics, otocs, and entanglement growth in systems without conservation laws, Phys. Rev. X 8, 021013 (2018).
- [10] T. Zhou and A. Nahum, Emergent statistical mechanics of entanglement in random unitary circuits, Phys. Rev. B 99, 174205 (2019).
- [11] T. Rakovszky, F. Pollmann, and C. W. von Keyserlingk, Diffusive hydrodynamics of out-of-time-ordered correlators with charge conservation, Phys. Rev. X 8, 031058 (2018).
- [12] V. Khemani, A. Vishwanath, and D. A. Huse, Operator spreading and the emergence of dissipative hydrodynamics under unitary evolution with conservation laws, Phys. Rev. X 8, 031057 (2018).
- [13] Y. Li, X. Chen, and M. P. A. Fisher, Quantum Zeno effect and the many-body entanglement transition, Phys. Rev. B 98, 205136 (2018).
- [14] B. Skinner, J. Ruhman, and A. Nahum, Measurement-induced phase transitions in the dynamics of entanglement, Phys. Rev. X 9, 031009 (2019).
- [15] S. Choi, Y. Bao, X. L. Qi, and E. Altman, Quantum error correction in scrambling dynamics and measurement-induced phase transition, Phys. Rev. Lett. **125**, 030505 (2020).
- [16] M. J. Gullans and D. A. Huse, Dynamical purification phase transition induced by quantum measurements, Phys. Rev. X 10, 041020 (2020).
- [17] Y. Li and M. P. A. Fisher, Statistical mechanics of quantum error correcting codes, Phys. Rev. B 103, 104306 (2021).
- [18] R. Fan, S. Vijay, A. Vishwanath, and Y.-Z. You, Self-organized error correction in random unitary circuits with measurement, Phys. Rev. B 103, 174309 (2021).
- [19] Y. Li, X. Chen, and M. P. A. Fisher, Measurement-driven entanglement transition in hybrid quantum circuits, Phys. Rev. B 100, 134306 (2019).
- [20] A. Chan, R. M. Nandkishore, M. Pretko, and G. Smith, Unitaryprojective entanglement dynamics, Phys. Rev. B 99, 224307 (2019).
- [21] Y. Li, X. Chen, A. W. W. Ludwig, and M. P. A. Fisher, Conformal invariance and quantum nonlocality in critical hybrid circuits, Phys. Rev. B 104, 104305 (2021).
- [22] X. Cao, A. Tilloy, and A. D. Luca, Entanglement in a fermion chain under continuous monitoring, SciPost Phys. 7, 024 (2019).

- [23] M. Szyniszewski, A. Romito, and H. Schomerus, Entanglement transition from variable-strength weak measurements, Phys. Rev. B 100, 064204 (2019).
- [24] Y. Bao, S. Choi, and E. Altman, Theory of the phase transition in random unitary circuits with measurements, Phys. Rev. B 101, 104301 (2020).
- [25] C.-M. Jian, Y.-Z. You, R. Vasseur, and A. W. W. Ludwig, Measurement-induced criticality in random quantum circuits, Phys. Rev. B 101, 104302 (2020).
- [26] M. J. Gullans and D. A. Huse, Scalable probes of measurementinduced criticality, Phys. Rev. Lett. 125, 070606 (2020).
- [27] A. Zabalo, M. J. Gullans, J. H. Wilson, S. Gopalakrishnan, D. A. Huse, and J. H. Pixley, Critical properties of the measurementinduced transition in random quantum circuits, Phys. Rev. B 101, 060301(R) (2020).
- [28] A. Nahum and B. Skinner, Entanglement and dynamics of diffusion-annihilation processes with Majorana defects, Phys. Rev. Res. 2, 023288 (2020).
- [29] M. Ippoliti, M. J. Gullans, S. Gopalakrishnan, D. A. Huse, and V. Khemani, Entanglement phase transitions in measurementonly dynamics, Phys. Rev. X 11, 011030 (2021).
- [30] A. Lavasani, Y. Alavirad, and M. Barkeshli, Measurementinduced topological entanglement transitions in symmetric random quantum circuits, Nat. Phys. 17, 342 (2021).
- [31] S. Sang and T. H. Hsieh, Measurement-protected quantum phases, Phys. Rev. Res. **3**, 023200 (2021).
- [32] Q. Tang and W. Zhu, Measurement-induced phase transition: A case study in the nonintegrable model by density-matrix renormalization group calculations, Phys. Rev. Res. 2, 013022 (2020).
- [33] J. Lopez-Piqueres, B. Ware, and R. Vasseur, Mean-field entanglement transitions in random tree tensor networks, Phys. Rev. B 102, 064202 (2020).
- [34] A. Nahum, S. Roy, B. Skinner, and J. Ruhman, Measurement and entanglement phase transitions in all-to-all quantum circuits, on quantum trees, and in Landau-Ginzburg theory, PRX Quantum 2, 010352 (2021).
- [35] X. Turkeshi, R. Fazio, and M. Dalmonte, Measurement-induced criticality in (2 + 1)-dimensional hybrid quantum circuits, Phys. Rev. B 102, 014315 (2020).
- [36] Y. Fuji and Y. Ashida, Measurement-induced quantum criticality under continuous monitoring, Phys. Rev. B 102, 054302 (2020).
- [37] O. Lunt, M. Szyniszewski, and A. Pal, Measurement-induced criticality and entanglement clusters: A study of 1D and 2D Clifford circuits, Phys. Rev. B 104, 155111 (2021).
- [38] O. Lunt and A. Pal, Measurement-induced entanglement transitions in many-body localized systems, Phys. Rev. Res. 2, 043072 (2020).
- [39] S. Vijay, Measurement-driven phase transition within a volumelaw entangled phase, arXiv:2005.03052.
- [40] X. Turkeshi, A. Biella, R. Fazio, M. Dalmonte, and M. Schiró, Measurement-induced entanglement transitions in the quantum Ising chain: From infinite to zero clicks, Phys. Rev. B 103, 224210 (2021).
- [41] M. Ippoliti and V. Khemani, Postselection-free entanglement dynamics via spacetime duality, Phys. Rev. Lett. 126, 060501 (2021).
- [42] T.-C. Lu and T. Grover, Entanglement transitions via space-time rotation of quantum circuits, PRX Quantum 2, 040319 (2021).

- [43] C.-M. Jian, B. Bauer, A. Keselman, and A. W. W. Ludwig, Criticality and entanglement in non-unitary quantum circuits and tensor networks of non-interacting fermions, arXiv:2012.04666.
- [44] S. Gopalakrishnan and M. J. Gullans, Entanglement and purification transitions in non-hermitian quantum mechanics, Phys. Rev. Lett. **126**, 170503 (2021).
- [45] X. Turkeshi, Measurement-induced criticality as a datastructure transition, Phys. Rev. B 106, 144313 (2022).
- [46] Y. Bao, S. Choi, and E. Altman, Symmetry enriched phases of quantum circuits, Ann. Phys. (NY) 435, 168618 (2021).
- [47] M. Block, Y. Bao, S. Choi, E. Altman, and N. Yao, The measurement-induced transition in long-range interacting quantum circuits, Phys. Rev. Lett. **128**, 010604 (2022).
- [48] G. Bentsen, S. Sahu, and B. Swingle, Measurement-induced purification in large-*N* hybrid Brownian circuits, Phys. Rev. B 104, 094304 (2021).
- [49] A. Zabalo, M. J. Gullans, J. H. Wilson, R. Vasseur, A. W. W. Ludwig, S. Gopalakrishnan, D. A. Huse, and J. H. Pixley, Operator scaling dimensions and multifractality at measurementinduced transitions, Phys. Rev. Lett. **128**, 050602 (2022).
- [50] U. Agrawal, A. Zabalo, K. Chen, J. H. Wilson, A. C. Potter, J. H. Pixley, S. Gopalakrishnan, and R. Vasseur, Entanglement and charge-sharpening transitions in U(1) symmetric monitored quantum circuits, Phys. Rev. X 12, 041002 (2022).
- [51] Y. Li and M. P. A. Fisher, Robust decoding in monitored dynamics of open quantum systems with \mathbb{Z}_2 symmetry, Phys. Rev. B 108, 214302 (2023).
- [52] O. Alberton, M. Buchhold, and S. Diehl, Entanglement transition in a monitored free-fermion chain: From extended criticality to area law, Phys. Rev. Lett. **126**, 170602 (2021).
- [53] S.-K. Jian, C. Liu, X. Chen, B. Swingle, and P. Zhang, Quantum error as an emergent magnetic field, arXiv:2106.09635.
- [54] M. Buchhold, Y. Minoguchi, A. Altland, and S. Diehl, Effective theory for the measurement-induced phase transition of Dirac fermions, Phys. Rev. X 11, 041004 (2021).
- [55] C. Noel, P. Niroula, D. Zhu, A. Risinger, L. Egan, D. Biswas, M. Cetina, A. V. Gorshkov, M. J. Gullans, D. A. Huse, and C. Monroe, Observation of measurement-induced quantum phases in a trapped-ion quantum computer, Nat. Phys. 18, 760 (2022).
- [56] R. Vasseur, A. C. Potter, Y.-Z. You, and A. W. W. Ludwig, Entanglement transitions from holographic random tensor networks, Phys. Rev. B 100, 134203 (2019).
- [57] R. Levy and B. K. Clark, Entanglement entropy transitions with random tensor networks, arXiv:2108.02225.
- [58] Z.-C. Yang, Y. Li, M. P. A. Fisher, and X. Chen, Entanglement phase transitions in random stabilizer tensor networks, Phys. Rev. B 105, 104306 (2022).
- [59] P. Hayden, S. Nezami, X.-L. Qi, N. Thomas, M. Walter, and Z. Yang, Holographic duality from random tensor networks, J. High Energy Phys. (online) 11 (2016) 009.
- [60] S. Nezami and M. Walter, Multipartite entanglement in stabilizer tensor networks, Phys. Rev. Lett. 125, 241602 (2020).
- [61] D. Gottesman, Class of quantum error-correcting codes saturating the quantum hamming bound, Phys. Rev. A 54, 1862 (1996).
- [62] D. Gottesman, The Heisenberg representation of quantum computers, arXiv:quant-ph/9807006.

- [63] S. Aaronson and D. Gottesman, Improved simulation of stabilizer circuits, Phys. Rev. A 70, 052328 (2004).
- [64] A. Chandran and C. R. Laumann, Semiclassical limit for the many-body localization transition, Phys. Rev. B 92, 024301 (2015).
- [65] D. Gross, S. Nezami, and M. Walter, Schur–Weyl duality for the Clifford group with applications: Property testing, a robust hudson theorem, and de Finetti representations, Commun. Math. Phys. 385, 1325 (2021).
- [66] D. Gottesman, Fault-tolerant quantum computation with higher-dimensional systems, Chaos, Solitons Fractals 10, 1749 (1999).
- [67] B. Collins, Moments and cumulants of polynomial random variables on unitary groups, the Itzykson-Zuber integral, and free probability, Int. Math. Res. Notices 2003, 953 (2003).
- [68] J. T. Chayes, L. Chayes, and R. Durrett, Critical behavior of the two-dimensional first passage time, J. Stat. Phys. 45, 933 (1986).
- [69] J. Jiang and C.-L. Yao, Critical first-passage percolation starting on the boundary, arXiv:1612.01803.
- [70] S. Sang, Y. Li, T. Zhou, X. Chen, T. H. Hsieh, and M. P. A. Fisher, Entanglement negativity at measurement-induced criticality, PRX Quantum 2, 030313 (2021).
- [71] M. Popp, F. Verstraete, M. A. Martín-Delgado, and J. I. Cirac, Localizable entanglement, Phys. Rev. A 71, 042306 (2005).
- [72] A. W. W. Ludwig, Infinite hierarchies of exponents in a diluted ferromagnet and their interpretation, Nucl. Phys. B 330, 639 (1990).
- [73] B. Duplantier and A. W. W. Ludwig, Multifractals, operatorproduct expansion, and field theory, Phys. Rev. Lett. 66, 247 (1991).
- [74] H. Apel, T. Kohler, and T. Cubitt, Holographic duality between local Hamiltonians from random tensor networks, J. High Energy Phys. (online) 03 (2022) 052.
- [75] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill, Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence, J. High Energy Phys. (online) 06 (2015) 149.
- [76] A. R. Calderbank and P. W. Shor, Good quantum errorcorrecting codes exist, Phys. Rev. A 54, 1098 (1996).
- [77] A. Steane, Multiple-particle interference and quantum error correction, Proc. R. Soc. London, Ser. A 452, 2551 (1996).
- [78] A. M. Steane, Error correcting codes in quantum theory, Phys. Rev. Lett. 77, 793 (1996).
- [79] D. Gross, Hudson's theorem for finite-dimensional quantum systems, J. Math. Phys. 47, 122107 (2006).
- [80] I. Roth, R. Kueng, S. Kimmel, Y.-K. Liu, D. Gross, J. Eisert, and M. Kliesch, Recovering quantum gates from few average gate fidelities, Phys. Rev. Lett. **121**, 170502 (2018).
- [81] B. Collins and P. Śniady, Integration with respect to the haar measure on unitary, orthogonal and symplectic group, Commun. Math. Phys. 264, 773 (2006).
- [82] B. Collins and S. Matsumoto, On some properties of orthogonal Weingarten functions, J. Math. Phys. 50, 113516 (2009).
- [83] B. Collins, S. Matsumoto, and J. Novak, The Weingarten calculus, Not. Am. Math. Soc. 69, 1 (2022).